On manifolds whose tangent bundle contains an ample locally free subsheaf

Marco Andreatta and Jarsoław A. Wiśniewski

Introduction.

Let X be a complex manifold of dimension n and let E be a vector bundle of rank r, or equivalently a locally free $\mathcal{O}_X$-sheaf of rank r. The bundle E is called ample if the relative hyperplane line bundle $O_{\mathbf{P}(E)}(1)$ over its projectivisation $\mathbf{P}(E) = Proj_X(Sym(E))$ is ample. We will assume that E is a subsheaf of the tangent sheaf TX, that is there exists a generically injective morphism $E \hookrightarrow TX$. In this paper we will prove the following:

Theorem. *If E is an ample locally free subsheaf of TX then $X \cong \mathbf{P}^n$ and $E \cong \mathcal{O}(1)^{\oplus r}$ or $E \cong T\mathbf{P}^n$.*

The characterization of $\mathbf{P}^n$ as the only manifold whose tangent bundle is ample was conjectured by R.Hartshorne. The Hartshorne conjecture was proved by S. Mori in a celebrated paper [Mo], which contained an amazing proof of the existence of rational curves on Fano manifolds. Building up on Mori's work a version of the present theorem was successively proved for $r = 1$ and $r = n,\ n-1,\ n-2$ by J. Wahl [Wa] and, respectively, F. Campana and T. Peternell [C-P]. Moreover Campana and Peternell posed a question with the above characterization of $\mathbf{P}^n$ which generalizes previous results.

The proof of the main theorem will apply rational curves on X. Our notation is consistent with the book of J. Kollár ([Ko]). In particular $Hom(\mathbf{P}^1, X)$ denotes the scheme parameterizing morphisms from $\mathbf{P}^1$ to X and $Hom(\mathbf{P}^1, X; 0 \to x)$ the scheme parameterizing morphisms sending $0 \in \mathbf{P}^1$ to $x \in X$. By $F : Hom(\mathbf{P}^1, X) \times \mathbf{P}^1 \to X$ we denote the evaluation morphism $F(f, p) = f(p)$. For any family of morphisms $V \subset Hom(\mathbf{P}^1, X)$ by F_V we denote the restriction of F to V and by $Locus(V)$ the closure of the image of F_V. We say that V is unsplit if the image of V in $Chow(X)$, via the natural morphism $[f] \mapsto [f_*(\mathbf{P}^1)]$, is proper.

If G is a vector bundle over $\mathbf{P}^1$ with G^+ we denote its positive part, that is the sub-bundle $[im(H^0(G(-1)) \otimes \mathcal{O} \to G(-1)] \otimes \mathcal{O}(1)$.

Prologue.

Let X and E be as in the introduction: we therefore assume that E is ample and moreover of rank $r > 1$, as the case $r = 1$ is set by Wahl's result.

By a theorem of Y. Miyaoka (see [Mi] or [Ko], IV.1.16) X is uniruled. So we can choose a closed irreducible component $V \subset Hom(\mathbf{P}^1, X)$ which covers X (that is $Locus(V) = X$) and which is a generically unsplit family (see [Ko], IV. 2.4). Moreover for a general $f \in V$ we have $f^*TX = \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus d} \oplus \mathcal{O}^{\oplus(n-d-1)}$ where $d = deg(f^*(-K_X)) - 2$, see ([Ko] IV 2.8, 2.9 and 2.10).

Lemma (0.1). *For any $f \in V$ the pull-back f^*E is isomorphic either to $\mathcal{O}(1)^{\oplus r}$ or to $\mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus(r-1)}$. In particular the family of curves parametrized by V is unsplit.*

Proof. For a general $f \in V$ the pull-back f^*E is an ample subbundle of $f^*TX = \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus(d)} \oplus \mathcal{O}^{\oplus(n-d-1)}$ and thus it is as in the lemma. Since E is ample this is true also for all $f \in V$. Since $deg(f^*E) = r$ or $deg(f^*E) = r + 1$ and $r > 1$, and for any ample bundle $\mathcal{E}$ over a rational curve we have $deg(\mathcal{E}) \geq rank(\mathcal{E})$, it follows that no curve from V can be split into a sum of two or more rational curves hence V is unsplit.

The family V defines a relation of *rational connectedness with respect to* V, which we shall call rc$V$ relation for short, in the following way: $x_1,\ x_2 \in X$ are in the rc$V$ relation if there exists a chain of rational curves parametrized by morphisms from V which joins x_1 and x_2. The rc$V$ relation is an equivalence relation and its equivalence classes can be parametrized generically by an algebraic set. More precisely, we have the following result due to Campana [Ca] and, independently, to Kollár-Miyaoka-Mori, [KMM2].

Theorem (0.2). *(see [Ko], IV.4.16). There exists an open subset $X^0 \subset X$ and a proper surjective morphisms with connected fibers $\varphi^0 : X^0 \to Z^0$ onto a normal variety, such that fibers of φ^0 are equivalence classes of rc$V$ relation.*

We shall call the morphism φ^0 an rc$V$ fibration. If Z_0 is just a point then we will call X a rationally connected manifold with the respect to the family V, in short an rc$V$ manifold. In what follows we shall analyze X using the notions of rc$V$ relation and rc$V$ fibration, the key is the following observation.

Lemma (0.3). *Let X, E and V be as above and moreover assume that $\varphi^0 : X^0 \to Z^0$ is an rc$V$ fibration. Then E is tangent to a general fiber of φ^0. That is, if X_g is a general fiber of φ^0, then the injection $E_{|X_g} \to TX_{|X_g}$ factors via $E_{|X_g} \hookrightarrow TX_g$.*

Proof. Choose a general X_g (in particular smooth) and let moreover $x \in X_g$ and $f \in V_x = V \cap Hom(\mathbf{P}^1, X; 0 \to x)$ be general as well. Then $Locus(V_x) \subset X_g$ and thus, by [Ko] II.3.4, $(f^*TX)^+_p \subset (f^*TX_g)_p$ for every $p \in \mathbf{P}^1 \setminus \{0\}$. This implies that $E_{|X_g} \to TX_{|X_g}$

factors to $E_{|X_g} \to TX_g$ generically and since the map $TX_g \to TX_{|X_g}$ has cokernel which is torsion free (it is the normal sheaf which is locally free) this yields $E_{|X_g} \hookrightarrow TX_g$, a sheaf injection.

Now the argument for the proof of the main theorem goes as follows. First we assume that X is a rc$V$ manifold and for $f \in V$ the pullback f^*E is either $\mathcal{O}(1)^{\oplus r}$ or $\mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus(r-1)}$ and we prove that $X \cong \mathbf{P}^n$ and $E \cong \mathcal{O}(1)^{\oplus r}$ or $E \cong T\mathbf{P}^n$, respectively. This is taken care of in the two subsequent sections: these two cases are reduced to Wahl's and Mori's theorems, respectively. In the last section we assume that the rc$V$ fibration is non-trivial: then we prove that it can be extended in codimension 1 so that we can produce a projective bundle $X_B \to B$ over a smooth curve B with an ample vector bundle $E_B \hookrightarrow T_{X_B/B}$. This is impossible, as observed by Campana and Peternell.

Trivial projective bundles over rc$V$ manifolds.

In this section we assume that X is a smooth projective manifold which is rationally connected with respect to an unsplit family V of rational curves (rc$V$ manifold). We let E be a rank r vector bundle. We begin by observing some general facts.

Proposition (1.1). *Let X be a projective manifold and $V \subset Hom(\mathbf{P}^1, X)$ an unsplit family of rational curves whose locus is the whole X. Then X is a rc$V$ manifold if and only if the Picard number $\rho(X)$ is 1.*

Proof. The "only if" part follows for instance from IV. 3.13.3 in [Ko] and the assumption on the unsplitteness of the family V. Note that this part is true even if X is not smooth. The "if" part (which will not be needed in the paper) is proved in [KMM1].

Let us note that rationally connectedness and $\rho(X) = 1$ imply that X is Fano, so in fact $PicX = \mathbf{Z}$. The following result concerns vector bundles of higher rank.

Proposition (1.2). *In the above notation suppose moreover that there exists an integer a such that for any $f \in V$ we have $f^*E = \mathcal{O}(a)^{\oplus r}$. Then there exists a (uniquely defined) line bundle L over X such that $deg f^*L = a$ and $E \cong L^{\oplus r}$.*

Proof. The argument applies induction with respect to r. Let us consider the projectivisation $p : \mathbf{P}(E) \to X$ with the relative $\mathcal{O}(1)$ bundle which we will denote by $\mathcal{L}$. We note that for any $f \in V$ and $y \in p^{-1}(f(0))$ we have a unique lift-up $\hat{f} : \mathbf{P}^1 \to \mathbf{P}(E)$ such that $p \circ \hat{f} = f$ and $deg(\hat{f}^*(\mathcal{L})) = a$, and $\hat{f}(0) = y$. That is, since $\mathbf{P}(f^*E) = \mathbf{P}^1 \times \mathbf{P}^{r-1}$, the morphism $\hat{f}$ is obtained by composing $\mathbf{P}(f^*E) \to \mathbf{P}(E)$ with the product morphism $\mathbf{P}^1 \to \mathbf{P}^1 \times \{y\} \subset \mathbf{P}^1 \times \mathbf{P}^{r-1}$. Thus, for a generic f we have $\hat{f}^*T\mathbf{P}(E) = f^*TX \oplus \mathcal{O}^{\oplus(r-1)}$. We can choose an irreducible $\hat{V} \subset Hom(\mathbf{P}^1, \mathbf{P}(E))$ which parameterizes these lift-ups, that is, via the natural morphism $p_* : Hom(\mathbf{P}^1, \mathbf{P}(E)) \to Hom(\mathbf{P}^1, X)$, defined by $p_*(\hat{f}) = p \circ \hat{f}$, the component $\hat{V}$ dominates V.

Claim (1.2.1). *The morphism $p_* : \hat{V} \to V$ is proper and thus surjective, and moreover $\hat{V}$ is an unsplit family.*

Proof. We use valuative criterion of properness [Hartshorne, II.4.7]. Let Δ be a spectrum of a discrete valuation ring (or a germ of a smooth curve in the analytic context) with a closed point δ and the general point Δ^0. Then for any family of morphims $F_\Delta : \Delta \times \mathbf{P}^1 \to X$ coming from $\Delta \to V$ we have $\mathbf{P}(F_\Delta^*(E)) = \Delta \times \mathbf{P}^1 \times \mathbf{P}^{r-1}$. Take now $\hat{F}_{\Delta_0} : \Delta^0 \times \mathbf{P}^1 \to \mathbf{P}(E)$, coming from a lift-up $\Delta^0 \to \hat{V}$ of $\Delta \to V$. By the construction $\hat{F}_{\Delta_0}$ is the composition of $\mathbf{P}(F_\Delta^*(E)) \to \mathbf{P}(E)$ with the product $id \times \psi_0 : \Delta_0 \times \mathbf{P}^1 \to (\Delta_0 \times \mathbf{P}^1) \times \mathbf{P}^{r-1}$, for some $\psi_0 : \Delta_0 \to \mathbf{P}^{r-1}$. The morphism ψ_0 extends to $\psi : \Delta \to \mathbf{P}^{r-1}$, thus $\hat{F}_{\Delta_0}$ extends to $\hat{F}_\Delta$ which is the composition of $\mathbf{P}(F_\Delta^*(E)) \to \mathbf{P}(E)$ with the product $id \times \psi$, hence p_* is proper.

The proof of unsplitting of $\hat{V}$ is similar. Namely let W and $\hat{W}$ denote the image of V and $\hat{V}$ in $Chow(X)$ and $Chow(\mathbf{P}(E))$, respectively. Let $p_* : \hat{W} \to W$ denote, by abuse of notation, the push-forward map. Similarly as above we prove that p_* is proper. Therefore $\hat{W}$ is proper and $\hat{V}$ is unsplit.

To proceed with the proof of the proposition let us consider the rc$\hat{V}$ fibration of $\mathbf{P}(E)$. Let $Y \subset \mathbf{P}(E)$ be a general fiber of this fibration. It is projective and smooth, and by the previous proposition $\rho(Y) = 1$. Moreover, by the surjectivity of $p_* : \hat{V} \to V$ and rational connectedness of X, the restriction map $p_Y : Y \to X$ is surjective and, since $\rho(Y) = 1$, it has no positive dimensional fiber, so it is a finite morphism. What's more, the restriction of $\mathcal{L}$ to Y, call it $\mathcal{L}_Y$, has intersection equal a with any curve from $\tilde{V} = \hat{V} \cap Hom(\mathbf{P}^1, Y)$.

Let us consider the pull-back $\tilde{p} : \mathbf{P}(p_Y^* E) \to Y$ with the induced morphism $\tilde{p}_Y : \mathbf{P}(p_Y^* E) \to \mathbf{P}(E)$ such that $p \circ \tilde{p}_Y = p_Y \circ \tilde{p}$. By the universal property of the fiber product the projective bundle $\tilde{p}$ admits a section $s : Y \to \mathbf{P}(p_Y^* E)$ such that $\tilde{p}_Y \circ s$ is the embedding of Y into $\mathbf{P}(E)$. This gives us a sequence of bundles over Y:

$$0 \longrightarrow E' \longrightarrow p_Y^*(E) \longrightarrow \mathcal{L}_Y \longrightarrow 0$$

where E' is a bundle of rank $r-1$ and over Y it satisfies the required assumptions with respect to the family $\tilde{V}$. Thus, by the inductive assumption, $E' \cong \mathcal{L}_Y^{\oplus(r-1)}$. But because Y is Fano, $H^1(Y, \mathcal{O}_Y) = 0$ and thus the above sequence of vector bundles splits; therefore $p_Y^*(E) \cong \mathcal{L}_Y^{\oplus r}$.

Now we shall be done by the following.

Lemma (1.2.2). *Let X be a Fano manifold with $p : \mathbf{P}(E) \to X$ a projectivisation of a rank r bundle. Suppose that $\Psi : Y \to X$ is a finite morphism. If $\mathbf{P}(\Psi^*(E)) \cong Y \times \mathbf{P}^{r-1}$ then $\mathbf{P}(E) \cong X \times \mathbf{P}^{r-1}$.*

Proof. By $\mathcal{L}$ let us denote the relative $\mathcal{O}(1)$ over $\mathbf{P}(E)$. We claim that $r\mathcal{L} - p^* detE$ is nef and $(r\mathcal{L} - p^* detE)^r = 0$ over $\mathbf{P}(E)$. This follows because the pull-back of $r\mathcal{L} - p^* detE$

to $\mathbf{P}(\Psi^*(E))$ has these features. By the same reason $r\mathcal{L} - p^*(detE + K_X)$ is ample and therefore $\mathbf{P}(E)$ is a Fano manifold and by Kawamata-Shokurov base-point-freeness $r\mathcal{L} - p^*detE$ defines a contraction $\varphi : \mathbf{P}(E) \to Z$ onto a normal projective variety of dimension $r-1$. Any fiber of φ is mapped, via p, surjectively onto X, with no positive dimensional fiber. Let T be a general fiber of φ. Then, T is smooth and by adjunction we find out that

$$K_T = (K_{\mathbf{P}(E)})_{|T} = (p^*K_X + (p^*(detE) - r\mathcal{L}))_{|T} = (p^*K_X)_{|T}$$

and therefore the restriction $p_{|T} : T \to X$ is unramified. Since X, being Fano, is simply-connected, it follows that T is a section of p. Thus we conclude that $Z \cong \mathbf{P}^{r-1}$ and $\mathbf{P}(E) \cong X \times \mathbf{P}^{r-1}$.

As an immediate consequence we get the following

Proposition (1.3). *Let X be a rationally connected manifold with respect to an unsplit family V and let E be an ample bundle which is a subsheaf of TX. If for $f \in V$ there is $deg(f^*E) = rank(E)$ then $X \cong \mathbf{P}^n$ and $E \cong \mathcal{O}(1)^{\oplus r}$.*

Proof. By the above splitting result we reduce the situation to the case $r = 1$, that is the Wahl's theorem (in the special case of $\rho(X) = 1$).

Tangent cone to a family of curves.

Let X be a smooth projective variety and $V \subset Hom(\mathbf{P}^1, X)$ and assume that V is generically unsplit so that for a general $[f] \in V$ we have $f^*TX = \mathcal{O}(2) \oplus \mathcal{O}(1)^{\oplus(d)} \oplus \mathcal{O}^{\oplus(n-d-1)}$; in particular for a general $[f]$ the map f is an immersion.

Fix a general $x \in X$ and consider $V_x = V \cap Hom(\mathbf{P}^1, X; 0 \mapsto x)$. Let t be a local coordinate around $0 \in \mathbf{P}^1$. Consider a derivative map $\Phi_x : V_x \to P(T_xX) = P((f^*TX)_0)$ which is defined at $[f] \in V_x$ if f is an immersion at 0 by $\Phi_x([f]) = [(Tf)_0(\partial/\partial t)]$, c.f. [Mori79, pp.602-603]. In the formula $Tf : T\mathbf{P}^1 \to f^*TX$ is the tangent map and T_xX is identified naturally, via f^*, with $(f^*TX)_0$.

By P we denote the "natural projectivisation" (that is vector spaces modulo homotheties) in opposition to "Grothendieck projectivisation" (that is projective spectrum of the symmetric algebra of a vector space) which we denote by $\mathbf{P}$. Using the latter formalism the map Φ_x is the value over 0 of the natural section of $\mathbf{P}(f^*\Omega_X)$ obtained by the surjective (at 0) morphism of derivatives: $Df : f^*\Omega_X \longrightarrow \Omega_{\mathbf{P}^1} \cong \mathcal{O}(-2)$.

We define $S_x \subset P(T_xX)$ as the closure of the image of the map Φ_x and we call it tangent cone of curves from V at the point x. J.-M.Hwang and N. Mok call this variety of minimal rational tangents [H-M]. The name of tangent cone follows from the fact that S_x is (at least around $[f]$) the tangent cone to $Locus(V_x)$. Indeed, let $\pi : \widehat{X_x} \to X$ be the blow-up of X at x with the exceptional divisor $E_x = P(T_xX)$. Consider $\hat{f}$:

$\mathbf{P}^1 \to \widehat{C} \subset \widehat{X_x}$, the lift-up of f, then $\hat{f}^*(T\widehat{X_x}) = \mathcal{O}(2) \oplus \mathcal{O}^{\oplus(d)} \oplus \mathcal{O}(-1)^{\oplus(n-d-1)}$. Thus $Hom(\mathbf{P}^1, \widehat{X_x})$ is smooth at $[\hat{f}]$ and of dimension $d+3$. Moreover, by [Ko] II.3.4 the evaluation morphism $\hat{F} : Hom(\mathbf{P}^1, \widehat{X_x}) \times \mathbf{P}^1 \to \widehat{X_x}$ is an immersion along $[\hat{f}] \times \mathbf{P}^1$ and moreover, by definition, $\hat{F}([\hat{f}], 0) = \Phi_x([f])$. On the other hand if we take an irreducible component $\hat{V}$ of $Hom(\mathbf{P}^1, \widehat{X_x})$ which contains $[\hat{f}]$ then $Locus(\hat{V})$ outside of E_x coincides with a component of $Locus(V_x)$. Thus around $\Phi_x([f])$ we get $S_x = E_x \cap \widetilde{Locus}(V_x)$, with $\widetilde{Locus}(V_x)$ denoting the strict transform of $Locus(V_x)$, so S_x is the tangent cone to $Locus(V_x)$.

For our purposes we need the following observation which follows from the above discussion (it can be find also in [Hw], proposition (2.3)).

Lemma (2.1). *The projectivised tangent space of the tangent cone S_x at $\Phi_x([f])$ is equal to $P((f^*TX)_0^+) \subset P((f^*TX)_0) = P(T_xX)$.*

Proof. By [Ko] II.3.4 the tangent space to $Locus(V_x)$ at $f(p)$ for $p \neq 0$ is the image of the evaluation of sections of the twisted pull-back of TX which is $im(T\hat{F})_p = (f^*TX)_p^+ \subset (f^*TX)_p = T_{f(p)}X$. Thus passing with p to 0 we get the result.

Lemma (2.2). *Let $V \subset Hom(\mathbf{P}^1, X)$ be as above and moreover suppose that $\mathcal{E} \hookrightarrow TX$ is a reflexive subsheaf with a torsionfree cokernel. If for a general $[f] \in V$ the tangent map $Tf : T\mathbf{P}^1 \to f^*TX$ factors to an injective $T\mathbf{P}^1 \hookrightarrow f^*\mathcal{E}$ then $(f^*TX)^+ \hookrightarrow f^*\mathcal{E}$.*

Proof. We choose a general f which is an immersion at $0 \to x$. Then $\Phi_x([f]) \in P(\mathcal{E}_x) = P((f^*\mathcal{E})_0) \subset P(T_xX) = P((f^*TX)_0)$ and the same holds for morphisms in a neighborhood of $[f]$ in V_x. Thus around $\Phi_x([f])$ the tangent cone S_x is contained in $P(\mathcal{E}_x) = P((f^*\mathcal{E})_0)$, so is its tangent space $P((f^*TX)_0^+)$.

Now we use it to conclude the case $f^*E \cong \mathcal{O}(2) \oplus \mathcal{O}(1)^{r-1}$ which we have singled out in our preliminary discussion. In view of Mori result we will be done by the following

Proposition (2.3). *Let X be a manifold rationally connected with respect to some unsplit $V \subset Hom(\mathbf{P}^1, X)$. Assume that E is an ample vector bundle admitting sheaf injection $E \to TX$ and for a general $[f] \in V$ we have $f^*E \cong \mathcal{O}(2) \oplus \mathcal{O}(1)^{r-1}$. Then $E \cong TX$.*

Proof. Comparing the splitting type of f^*E and f^*TX we see that the tangent map $Tf : f^*T\mathbf{P}^1 \to f^*TX$ factors to a vector bundle (nowhere degenerate) injection $T\mathbf{P}^1 \to f^*E$. In other words, we have surjective morphism $(f^*E)^* \to \Omega_{\mathbf{P}^1} \cong \mathcal{O}(-2)$. Thus the values of $\Phi_{f(p)}$ at any point $p \in \mathbf{P}^1$ are in $P(E)$. This holds also for small deformations of the morphism f at any point $p \in \mathbf{P}^1$ and therefore (the component of) tangent cone $S_{f(p)}$ is contained in $P(E_{f(p)})$. Thus, in view of the previous lemma, we have the inclusion $(f^*TX)^+ \subset f^*E$ and hence, by the splitting type of f^*TX, we conclude that $f^*E =$

$(f^*TX)^+$ and therefore $deg(f^*E) = deg(f^*(-K_X))$. Since $\rho(X) = 1$ it follows that $det(E) = -K_X$.

The embedding $E \hookrightarrow TX$ gives rise to a non-trivial morphism $det(E) \to \Lambda^r TX$ and thus to a non-zero section of $\Lambda^r TX \otimes K_X$. We use dualities to have the equalities:

$$h^0(X, \Lambda^r TX \otimes K_X) = h^n(X, \Omega^r_X) = h^r(X, \Omega^n_X) = h^r(X, K_X) = h^{n-r}(X, \mathcal{O}_X)$$

and, since X is Fano, the latter number is non-zero only if $r = n$. Thus $\Lambda^r TX \otimes (detE)^{-1} \cong \mathcal{O}_X$ so $E \hookrightarrow TX$ is nowhere degenerate, hence an isomorphism.

Extending rc$V$ fibrations in codimension 1.

Let X, E be as in the main theorem with the unsplit family V and $\varphi^0 : X^0 \to Z^0$ be the rc$V$ fibration as defined at the beginning of the paper. In this section we assume that $dimZ^0 \geq 1$ which will lead to a contradiction.

A general fiber of φ^0, call it X_g, is rationally connected and, as we have noticed, E is tangent to X_g , that is the injection $E_X \to TX_{X_g}$ factors to $E_{X_g} \to TX_g$.

By the result of the previous sections X_g is isomorphic to $\mathbf{P}^k$ and E_{X_g} is either $TX_g = T\mathbf{P}^k$ or $\mathcal{O}(1)^{\oplus r}$. We can shrink Z^0 and X^0 so that $\varphi^0 : X^0 \to Z^0$ is a projective bundle in étale or analytic topology.

Take now $\tilde{Z}$ an irreducible component of $Hilb(X)$ which contains a general fiber X_g. Over $\tilde{Z}$ there exists a universal flat family $\tilde{X} \subset \tilde{Z} \times X$ with projections $\tilde{\varphi} : \tilde{X} \to \tilde{Z}$ and $\tilde{\beta} : \tilde{X} \to X$. The family $\tilde{X} \to \tilde{Z}$ extends $X^0 \to Z^0$, that is we have an inclusion $X^0 \hookrightarrow \tilde{X}$ and $Z^0 \hookrightarrow \tilde{Z}$ such that $\tilde{\varphi}$ extends φ^0.

Lemma (3.1). *For any $z \in Z$ the fiber $\tilde{X}_z := \tilde{\varphi}^{-1}(z)$ is irreducible.*

Proof. This follows from the fact that any two points in $\tilde{X}_z$ can be joint by an irreducible curve parametrized by a morphism from V. Indeed, take x_1, $x_2 \in \tilde{X}_z$, then there exists a 1-parameter family of $X_t \cong \mathbf{P}^k$ whose limit is $\tilde{X}_z$ with points x_1^t, $x_2^t \in X_t$ whose limits are x_1 and x_2, respectively. Now x_1^t can be joint with x_2^t by a line C_t parametrized by a $[f_t] \in V$. Since V is unsplit the limit curve is irreducible and the claim follows.

The morphism $\tilde{\beta}$ is birational with exceptional locus $E(\tilde{\beta})$ which is of codimension 1, by Zariski main theorem. If we set $X^* = \tilde{X} \setminus E(\tilde{\beta})$ then, via $\tilde{\beta}$, we have inclusion $X^* \subset X$ and $X \setminus X^*$ is of codimension ≥ 2. Let φ^* be the restriction of $\tilde{\varphi}$ to X^* and let $Z^* = \varphi^*(X^*)$.

Lemma (3.2). *The morphism $\varphi^* : X^* \to Z^*$ is proper.*

Proof. Since $\tilde{\varphi}$ is proper, it is enough to show that if $E(\beta)$ meets a fiber of $\tilde{\varphi}$ then it contains all of it. Let $E(\beta) = \bigcup E_i$ be a decomposition into irreducible components. Since

$\tilde{\varphi}$ extends φ^0 defined on a subset of X it follows that none of E_i meets a generic fiber of $\tilde{\varphi}$ so $dim(\tilde{\varphi}(E_i)) \leq dimZ - 1 = n - k - 1$. As dimension of any fiber of $\tilde{\varphi}_{|E_i}$ is $\leq k$ it follows that, actually, all fibers of $\tilde{\varphi}_{|E_i}$ are of dimension k. But all fibers of $\tilde{\varphi}$ are irreducible and of dimension k hence our claim.

Let $\hat{\varphi} : \hat{X} \to \hat{Z}$ be the normalization of $\tilde{\varphi} : \tilde{X} \to \tilde{Z}$, with the induced morphism $\hat{\beta} : \hat{X} \to X$. Since $X^* \subset \tilde{X}$ is smooth it lifts up to $X^* \subset \hat{X}$. The restriction $\hat{\varphi}$ to X^*, call it $\hat{\varphi}^*$, is proper and call its image $\hat{Z}^*$.

Lemma (3.4). *Outside a subset of codimension ≥ 2 the morphism $\hat{\varphi}^*$ is a $\mathbf{P}^k$-bundle (in the analytic topology).*

Proof. This is a result of Fujita. Let B be a curve obtained by intersection of $dimZ - 1$ general very ample divisor on $\hat{Z}$ and let $B^* = B \cap \hat{Z}^*$. By Bertini B^* is smooth and $X^*_B = (\hat{\varphi}^*)^{-1}(B) \cap X^*$ is smooth as well. Moreover, the induced morphism $\hat{\varphi}_B : X^*_B \to B^*$ is generically projective bundle, so it is a projective bundle by [Fu], lemma (2.12). Thus, by Bertini, outside of codimension ≥ 2 the morphism $\hat{\varphi}^*$ is smooth and with fibers equal to $\mathbf{P}^k$.

Let $\varphi' : X' \to Z'$ be the restriction of $\hat{\varphi}$ which is a $\mathbf{P}^k$-bundle. By the above lemmas codimension of $X \setminus X'$ is ≥ 2 and therefore we can take a general smooth projective curve B' in X which is contained in X' and is not contained in fibers of φ'. Then $B = \varphi'(B')$ is projective and contained in Z', and since our choice was general it is moreover smooth. Let $X_B = (\varphi')^{-1}(B)$. Again, by the generality of the choice we can assume that $E_{|X_B} \to TX_{|X_B}$ is a subsheaf inclusion which is generically of maximal rank.

Lemma (3.5). *In the above notation $E_{|X_B}$ is a locally free subsheaf of $T_{X_B/B}$.*

Proof. We have noticed at our initial argument, that the vector bundle E is generically tangent to fibers of φ^0. Therefore the sheaf inclusion $E_{|X_B} \hookrightarrow TX_{|X_B}$ generically factors to $E_{|X_B} \to T_{X_B/B}$. Since the cokernel of the composition $T_{X_B/B} \to TX_B \to TX_{|X_B}$ is torsion free (it is actually locally free) the inclusion $E_{|X_B} \hookrightarrow T_{X_B/B}$ over X_B follows.

Conclusion. We arrive to a contradiction by applying Proposition 4.5 of [C-P] to the $\mathbf{P}^k$ bundle $f^* : X_B \to B$ and to the ample vector bundle $E_{|X_B}$.

Acknowledgments.

The first named author was partially supported by the MURST of the Italian Government. Some ideas of this paper was conceived while the second named author had Alexander von Humboldt fellowship in Göttingen, Germany; he would like moreover to acknowledge support of Polish KBN (2P03A02216) and to thank Universitá di Trento, Italy, for hospitality.

References.

[Ca] Campana, F., Connexite rationelle des varieties de Fano, Ann. Sci. Ec. Norm. Sup. 25, 539-545 (1992).

[C-P] Campana, F., Peternell, Th., Rational curves and ampleness properties of tangent bundle of algebraic varieties, Manuscripta Math., **97** (1998), 59-74.

[Fu] Fujita, T., On polarized manifolds whose adjoint bundles are not semipositive, in *Algebraic Geometry, Sendai 1985*, Adv. Stud. Pure Math., **10** (1987), 167-178.

[Ha] Hartshorne,R., *Algebraic geometry*, GTM **52** (1977), Springer-Verlag.

[HM] Hwang, J.-M., Mok, N., Holomorphic maps from rational homogeneous spaces of Picard number 1 onto projective manifolds, Inventiones Math. 136, 208-236 (1999).

[Hw] Hwang, J.-M., Geometry of minimal rational curves on Fano manifolds, preprint of a lecture given at the School on Vanishing Theorem etc., Trieste 2000.

[Ko] Kollár, J., *Rational Curves on Algebraic Varieties*, Ergebnisse der Math. **32** (1995), Springer Verlag.

[KMM1] Kollár, J., Miyaoka, Y., Mori, Sh., Rational curves on Fano varieties, in *Trento 1990*, Lecture Notes in Math 1515.

[KMM2] Kollár, J., Miyaoka, Y., Mori, Sh., Rationally connected varieties, J. Alg. Geom. 429-448 (1992).

[Mi] Miyaoka, Y., The Chern classes and Kodaira dimension of a minimal variety, in *Algebraic Geometry, Sendai 1985*, Adv. Stud. Pure Math., **10** (1987), 449-476.

[Mo] Mori, Sh., Projective manifolds with ample tangent bundles, Ann. Math. **110** (1979), 593-606.

[Wa] Wahl, J., A cohomological characterization of P^n, Inv. Math. 72 315-322 (1983).

Marco Andreatta, Dipartimento di Matematica, Università di Trento, 38050 Povo, Italy
e-mail: andreatt@science.unitn.it
Jarosław A. Wiśniewski, Instytut Matematyki UW, Banacha 2, 02-097 Warszawa, Poland
e-mail: jarekw@mimuw.edu.pl