

# Complex Manifolds 2022/23

Lecture summary. **This is not a replacement for a textbook.**

aweber at mimuw.edu.pl

<http://www.mimuw.edu.pl/~aweber/ComplexManifolds>

Date August 21, 2023

The main reference: Daniel Huybrechts, Complex Geometry, An introduction. (Springer 2005)

Also:

Donu Arapura, Algebraic Geometry over the Complex Numbers (Universitext)

Claire Voisin, Hodge Theory and Complex Algebraic (Cambridge Studies in Advanced Mathematics)

## 1 Introduction

### 1.1 Definition of complex manifolds

### 1.2 Projective spaces

### 1.3 Grassmannians $Gr_k(\mathbb{C}^n)$ . Affine maps: for $I = \{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$

$$U_I = \{V \in Gr_k(\mathbb{C}^n) \mid \text{projection } V \rightarrow I\text{-coordinates is an isomorphism}\}$$

$$U_I \simeq \text{Hom}(\mathbb{C}^k, \mathbb{C}^{n-k}).$$

### 1.4 Plücker embedding, $Gr_2(\mathbb{C}^4)$ as a quadric in $\mathbb{P}^5 = \mathbb{P}(\Lambda^2 \mathbb{C}^4)$ .

### 1.5 Hyperplane in $\mathbb{P}^n$ e.g. elliptic curve in $\mathbb{P}^2$

$$y^3 + pxz^2 + qz^3 - x^2z = 0$$

with  $p, q$  fixed.

**1.6** Complex manifolds as real manifolds are orientable since any linear complex map preserves the distinguished orientation of the underlying real vector space.

**1.7** Basic information about topological coverings and induced complex structures: If  $f : X \rightarrow Y$  is a topological covering,  $Y$  has a structure of a complex manifold, then  $X$  has a natural structure of a complex manifold and  $f$  is a holomorphic.

## Curves

**1.8** Riemann surfaces (= oriented surfaces with a Riemannian metric) and complex surfaces: each Riemannian surface has a complex structure. Genus of Riemann surface.

- The rotation by  $90^\circ$  in the tangent space allows to introduce a structure of complex vector space. This structure is „integrable” i.e. it comes from a structure of a complex manifold (a proof will be later, it follows trivially from Newlander-Nirenberg theorem).

**1.9** Riemann uniformization theorem: any complex curve is isomorphic to  $\mathbb{P}^1$  or it is a quotient of  $\mathbb{C}$  or  $\mathbb{D} \simeq \mathbb{H}$ .

- Another formulation: any simply connected complex curve is isomorphic to  $\mathbb{P}^1$ ,  $\mathbb{C}$  or  $\mathbb{D}$ . This is a generalization of the Riemann theorem for open subsets in  $\mathbb{C}$ .

**1.10** The automorphism group of  $\mathbb{P}^1$  is equal to  $PGL_2(\mathbb{C})$ . Any complex-analytic automorphism of  $\mathbb{P}^1$  is given by a linear formula. (The same statement holds for  $\mathbb{P}^n$ .)

- Proof: Composing with a linear map we can assume that  $f(0) = 0$ ,  $f(\infty) = \infty$ . Expanding at infinity we get an estimation  $1/|f(z)| < c/|z|$ . Hence the function  $g(z) = z/f(z)$  is bounded. It has no

poles, since at 0 the zero of the denominator cancels out and there are no more zeros of  $f$ . Hence by Liouville theorem  $g(z)$  is constant.

- Hence each automorphism of  $\mathbb{P}^1$  has a fixed point – the eigenvector of the linear map.
- Topological proof: there are no nontrivial topological covering  $\mathbb{P}^1 \simeq S^2 \rightarrow C$  except  $C = \mathbb{RP}^2$ . But the real projective plane is not orientable, so it cannot be a complex curve.

**1.11** Automorphisms of  $\mathbb{C}$  are given by affine maps  $f(z) = az + b$ . There are no fixed points only if  $a = 1$ .

- The map  $f : \mathbb{C} \rightarrow \mathbb{C}$  extends to  $\mathbb{P}^1$ . It is continuous at  $\infty$ . By Riemann extension theorem it is holomorphic at  $\infty$  and we apply (1.10).

**1.12** The complex quotients of  $\mathbb{C}$  are of the form  $\mathbb{C}/\Lambda$  for a lattice  $\Lambda \subset \mathbb{C}$ .

- The nontrivial discrete subgroups of  $\Lambda \subset (\mathbb{C}, +) \simeq \mathbb{R}^2$  are of the form  $\Lambda = \langle a, b \rangle$  for  $b/a \in \mathbb{H}$ , (or  $\Lambda = \langle a \rangle$ ). We can restrict our attention to subgroups of the form  $\Lambda = \langle 1, \tau \rangle$ ,  $\tau \in \mathbb{H}$ .
- The group  $PSL_2(\mathbb{C}) := SL_2(\mathbb{C})/\{\pm I\}$  acts on  $\mathbb{P}^1$  by homography:  $\begin{pmatrix} s & t \\ u & v \end{pmatrix} \cdot z = (sz + t)/(uz + v)$ . The subgroup  $PSL_2(\mathbb{R})$  preserves the upper hyperplane  $\mathbb{H}$ .
- Suppose  $\tau, \tau' \in \mathbb{H}$ . Then  $\mathbb{C}/\langle 1, \tau \rangle \simeq \mathbb{C}/\langle 1, \tau' \rangle$  if and only if  $\tau$  and  $\tau'$  belong to the same orbit of  $PSL_2(\mathbb{Z})$ . (Exercise)

**1.13** The group of disk automorphisms is isomorphic to the group of the upper hyperplane  $\mathbb{H}$  automorphisms:  $Aut(\mathbb{H}) = PSL_2(\mathbb{R})$ .

- $Aut(\mathbb{D})$  consist of homographies (apply the Schwartz lemma, assuming  $f(0) = 0$ ).

**1.14** Discrete subgroups of  $PSL_2(\mathbb{R})$  are called Fuchsian groups (grupy Fiksa). The curves of higher genera  $g > 1$  are quotients  $\mathbb{H}/G$  where  $G \subset PSL_2(\mathbb{R})$  is Fuchsian and acts without fixed points.

**1.15** Read more: [Huybrechts, Complex Geometry, Chapter 2.1]

## 2 Weierstrass preparation

Local theory: see [§1, Huybrechts].

**2.1** Cauchy-Riemann operator  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + i \frac{\partial}{\partial y}$  and complex differential  $\frac{1}{2} \frac{\partial}{\partial x} - i \frac{\partial}{\partial y}$ .

**2.2** Differentials  $dz = dx + idy$  and  $d\bar{z} = dx - idy$ .

- For any  $C^\infty$  function on  $f : \mathbb{C} \rightarrow \mathbb{C}$  the differential

$$df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy = \frac{\partial f}{\partial z} dz + \frac{\partial f}{\partial \bar{z}} d\bar{z}.$$

(Hint if  $A = \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix}$  then  $(A^T)^{-1} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix}$ .)

**2.3** Recollection of theorems for complex analytic functions in one variable

- series expansion
- Cauchy integration formula
- maximum principle
- identity principle
- Liouville theorem

**2.4** Residue  $res_{z_0}(f) = \frac{1}{2\pi i} \int_{\partial D_{z_0}} f dz$ , where  $D_{z_0}$  is a small disk around  $z$ .

**2.5** Residue theorem: for a meromorphic function  $f$  (enough to assume: holomorphic away from a discrete set  $\{z_1, z_2, \dots, z_n\}$ ) on a compact Riemann surface  $S$

$$\sum_k \operatorname{res}_{z_k}(f) = 0.$$

- Proof from the Stokes theorem: Assume that the discs  $D_{z_k}$  for  $z \in \operatorname{Sing}(f)$  do not intersect:

$$\sum_{z_k} \int_{\partial D_{z_k}} f dz = - \int_{\partial(S \setminus \bigcup D_{z_k})} f dz = - \int_{S \setminus \bigcup D_{z_k}} d(f dz) = - \int_{S \setminus \bigcup D_{z_k}} \frac{\partial f}{\partial \bar{z}} d\bar{z} \wedge dz = 0.$$

**2.6** A formula for the number of zeros in a disk has a generalization which will be used later. If  $f(z) \neq 0$  for  $|z| = \varepsilon$  then for  $\ell \geq 0$  we have

$$\frac{1}{2\pi i} \int_{S_\varepsilon} \frac{f'(\xi)}{f(\xi)} \xi^\ell d\xi = \sum_{|\alpha| < \varepsilon, f(\alpha)=0} \alpha^\ell.$$

## Many variables - references to [Huybrechts §1.1]

**2.7** Definition: a  $C^\infty$  function  $f : \mathbb{C}^n \rightarrow \mathbb{C}$  is holomorphic if  $\partial_{\bar{z}_k} f = 0$  for  $k = 1, 2, \dots, n$ .

**2.8** Cauchy integral formula Prop 1.1.2

**2.9** Hartogs theorem Prop 1.1.4

**2.10** Corollary: zero set of a holomorphic function ( $f \not\equiv 0$ ) has real codimension equal 2 or it is empty.

- Remark: any analytic set (eg zero set of a holomorphic function) is triangulable by Łojasiewicz theorem, so there is no ambiguity with the notion of dimension.

**2.11** Weierstrass preparation theorem (Th. 1.1.6).

**2.12** Algebraic fact used in the proof: elementary symmetric functions  $\sigma_k$  can be expressed by power sums  $p_k$ .

## Local ring

**2.13** The local ring  $\mathcal{O}_{\mathbb{C}^n,0}$  is a unique factorization domain (Prop 1.1.15).

- Key argument: Weierstrass polynomial is indecomposable in  $\mathcal{O}_{\mathbb{C}^{n-1},0}[z]$  iff it is indecomposable in  $\mathcal{O}_{\mathbb{C}^n,0}$ .

## 3 Weierstrass II

**3.1** Weierstrass preparation theorem – division version (Prop 1.1.17).

**3.2** The local ring  $\mathcal{O}_{\mathbb{C}^n,0}$  is noetherian (Prop 1.1.18).

**3.3** Remark: If  $\emptyset \neq U \subset \mathbb{C}^n$ ,  $n > 0$  then  $\mathcal{O}_{\mathbb{C}^n}(U)$  is not noetherian.

- Proof: Any  $I \subset \mathcal{O}_{\mathbb{C}^n,0}$  is generated by  $I \cap (\mathcal{O}_{\mathbb{C}^{n-1},0}[z])$  and any Weierstrass polynomial  $g \in I$  (by division version of WPT).

**3.4** Germ of sets and ideals in the local ring:

- The germ of the set  $Z(J)$  defined by an ideal  $J \subset \mathcal{O}_{\mathbb{C}^n,0}$ .  
— if  $J_1 \subset J_2$  then  $Z(J_1) \supset Z(J_2)$
- The ideal of function germs vanishing on the germ of a set  $I(X)$ . We have:  
— if  $X_1 \subset X_2$  then  $I(X_1) \supset I(X_2)$

### 3.5 Compositions of $Z$ and $I$

- $X \subset Z(I(X))$  for any set germ,
- $J \subset I(Z(J))$  for any ideal,
- $X = Z(I(X))$  for analytic set germs (i.e. of the form  $X = Z(J)$ )  
— since  $J \subset I(Z(J))$  then  $X = Z(J) \supset Z(I(Z(J))) = Z(I(X))$ .
- Hilbert nullstellensatz:  $I(Z(J)) = \sqrt{J}$  (see sketch of a proof in Huybrechts p.20).

**3.6** Let  $g \in \mathcal{O}_{\mathbb{C}^n,0}$  be indecomposable, then if  $f|_{Z(g)} = 0$ , then  $g$  divides  $f$  (Cor. 1.1.9)

- Proof from the division version of Weierstrass preparation theorem.
- Key step: if  $g$  is indecomposable Weierstrass polynomial, then  $g_w(z)$  generically (w/r to  $w$ ) has distinct roots.  
— let  $K$  be the quotient field of  $\mathcal{O}_{\mathbb{C}^{n-1},0}$ . The polynomials  $g_w(z)$  and  $g'_w(z)$  are coprime (by Gauss lemma), so there exist  $\alpha(z), \beta(z) \in K(z)$  such that  $\alpha(z)g_w(z) + \beta(z)g'_w(z) = 1$ . Passing to  $\mathcal{O}_{\mathbb{C}^{n-1},0}$ , removing the denominators

$$\tilde{\alpha}(z)g_w(z) + \tilde{\beta}(z)g'_w(z) = \gamma$$

with  $0 \neq \gamma \in \mathcal{O}_{\mathbb{C}^{n-1},0}$ . At the points where  $\gamma(w) \neq 0$  the polynomial  $g_w$  does not have multiple roots.

**3.7** The germ of a set is indecomposable (also called irreducible) if and only if  $I(X)$  is a prime ideal (Lemma 1.1.28)

**Rough notes on GAGA** (dla absolwentów teorii snopów)

*J.-P. Serre, Géométrie algébrique et géométrie analytique, Ann. Inst. Fourier 6: 1-42, (1956)*

See also: Amnon Neeman, Algebraic and analytic geometry. Cambridge University Press (2007)

**3.8** For an algebraic manifold  $X$  (a scheme in general) we define „analytification”  $X^{an}$ .

- As a set  $X = X^{an}$ .
- While  $X$  has Zariski topology,  $X^{an}$  has classical topology (glued from the open subsets  $U \subset \mathbb{C}^n \simeq \mathbb{R}^{2n}$ ). The identity map  $\iota : X^{an} \rightarrow X$  is continuous (every Zariski open set is open in the classical topology). [Serre §5 Lemma 1]
- Both spaces are ringed. We have distinguished sheaves of rings  $\mathcal{O}_X$  (algebraic functions) and  $\mathcal{H}_X$  (holomorphic functions), the stalks are local rings. We have a map

$$\theta_X : \iota^{-1}\mathcal{O}_X \rightarrow \mathcal{H}_X,$$

i.e.  $\iota$  extends to a map of ringed spaces. Here  $\iota^{-1}$  denotes the pull-back of a sheaf. The map  $\theta_X$  is injective, flat, an isomorphism after completion in  $\mathfrak{m}$ . [Serre §6, prop 4]

**3.9** For an algebraic sheaf  $\mathcal{F}$  over an algebraic manifold we define „analytification”

$$\mathcal{F}^{an} = \mathcal{H}_X \otimes_{\iota^{-1}\mathcal{O}_X} \iota^{-1}\mathcal{F}.$$

Of course  $\mathcal{O}_X^{an} = \mathcal{H}_X$ . [Serre §9, Prop 10]

**3.10** Definition: Let  $(Y, \mathcal{R}_Y)$  be a ringed space. The sheaf  $\mathcal{R}_Y$ -modules  $\mathcal{F}$  is coherent iff

- 1) locally there is a surjective map  $(\mathcal{R}_Y^N)|_U \rightarrow \mathcal{F}|_U$  for some  $N$  (i.e.  $\mathcal{F}$  is locally finitely generated),
- 2) for any map  $(\mathcal{R}_Y^M)|_U \rightarrow \mathcal{F}|_U$  the kernel is finitely generated.

By  $Coh(Y)$  we denote the category of coherent sheaves.

- Mind the difference comparing with the definition for algebraic varieties.

**3.11** Oka Theorem:  $\mathcal{F} = \mathcal{H}_X$  is coherent. (This is not a tautology!) References in [Serre §3 Prop.1]

**3.12** Analytification of sheaves is a functor preserving coherent sheaves [Serre §9]

$$(-)^{an} : Sh(X) \rightarrow Sh(X^{an})$$

**3.13** (Serre) If  $X$  is projective,  $\mathcal{F}$  coherent then the natural map  $H^*(X; \mathcal{F}) \rightarrow H^*(X^{an}; \mathcal{F}^{an})$  is an isomorphism. [Serre §12 Th. 1]

• Relative version: Let  $f : X \rightarrow Y$  be a projective morphism of algebraic varieties. Then  $f$  induces a functor

$$f_*^{an} : Coh(X^{an}) \rightarrow Coh(Y^{an})$$

and

$$\begin{aligned} (f_* \mathcal{F})^{an} &= f_*^{an} \mathcal{F}^{an} \\ (R^k f_* \mathcal{F})^{an} &= R^k f_*^{an} \mathcal{F}^{an} \end{aligned}$$

If  $Y = pt$  then we recover the previous formulation.

**3.14** (Serre cont.) If  $X$  is a projective variety, then  $(-)^{an}$  restricted to  $Coh(X)$  is an equivalence of categories.

The above means:

(i)  $Hom_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \rightarrow Hom_{\mathcal{H}_X}(\mathcal{F}^{an}, \mathcal{G}^{an})$  is an isomorphism. [Serre §12 Th. 2]

(ii) For any analytic coherent sheaf  $G$  there exists an algebraic sheaf  $\mathcal{F}$  such that  $G \simeq \mathcal{F}^{an}$ . [Serre §12 Th. 3]

**3.15** The proofs can be reduced to  $X = \mathbb{P}^n$ . To check the equality  $H^*(X; \mathcal{F}) \simeq H^*(X^{an}; \mathcal{F}^{an})$  we can assume (by various cohomology exact sequences) that  $\mathcal{F} \simeq \mathcal{O}(m)$ .

**3.16** For a proof of (i) use the equality of sheaf-Homs

$$(\underline{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}))^{an} = \underline{Hom}_{\mathcal{H}_X}(\mathcal{F}^{an}, \mathcal{G}^{an})$$

which holds for algebraic coherent sheaves. Then apply the general principle

$$Hom_Y(F, G) = H^0(Y; \underline{Hom}_Y(F, G)),$$

and apply 3.13.

**3.17** For a proof of (ii) have to show that any analytic sheaf  $F$  on  $X = \mathbb{P}^n$  after tensoring with  $\mathcal{H}_X(m)$  for some big  $m$  is globally generated, i.e. there exists  $k$  and a surjection

$$\mathcal{H}_X^k \rightarrow F(m) := F \otimes_{\mathcal{H}_X} \mathcal{H}_X(m),$$

which is equivalent to: for each point  $x \in X$

$$\text{Global sections of } F(m) \rightarrow F(m)_x$$

is a surjection, [Serre §16 Lemma 8]. Then  $F(m) = \text{coker}(\mathcal{H}_X^\ell \rightarrow \mathcal{H}_X^k)$ , thus by (i) it is algebraic, [Serre §17].

**3.18** Corollary (Chow Theorem): Any analytic subvariety  $\mathbb{P}^n$  is described by a set of polynomial equations..

## 4 Morse theory for $C^\infty$ -manifolds and weak Lefschetz

[Milnor – Morse theory, 1963]

**4.1** Def: Morse function  $f : M \rightarrow \mathbb{R}$  is a proper smooth function such that if  $Df(p) = 0$  for  $p \in M$  then  $D^2f(p)$  is nondegenerate. Additionally we assume that for each critical value there exist only one critical point of  $f$  (critical values of distinct points do not collide).

**4.2**  $ind(p)$  = the index of a critical point = the number of minuses after diagonalization of  $D^2f(p)$ .

4.3 for  $t \in \mathbb{R}$  let

$$M_{\leq t} = \{p \in M \mid f(p) \leq t\}.$$

4.4 Theorem:

- 1) If there is no critical value in the interval  $[a, b]$ , then the inclusion  $M_{\leq a} \subset M_{\leq b}$  is a homotopy equivalence
- 2) If  $f(p) = c \in [a, b]$  is the only one critical value in the interval  $[a, b]$  then  $M_{\leq b}$  is homeomorphic to  $M_{\leq a}$  with attached  $I^{ind(p)} \times I^{n-ind(p)}$  along  $\partial I^{ind(p)} \times I^{n-ind(p)}$ , (up to homotopy we attach a cell of the dimension  $k = ind(p)$ ).

4.5 The effect of attaching  $k$ -dimensional cell:

$$M_{\leq b} = M_{\leq a} \cup_{\phi} D^k, \quad \phi : S^{k-1} \rightarrow M_{\leq a}.$$

There is an exact sequence

$$\begin{array}{ccccccccccc} 0 & \rightarrow & H^{k-1}(M_{\leq b}) & \rightarrow & H^{k-1}(M_{\leq a}) & \xrightarrow{\phi^*} & H^{k-1}(S^{k-1}) & \rightarrow & H^k(M_{\leq b}) & \rightarrow & H^k(M_{\leq a}) & \rightarrow & 0 \\ & & & & & & \parallel & & & & & & \\ & & & & & & \mathbb{Z} & & & & & & \end{array}$$

The for the remaining gradations  $H^i(M_{\leq b}) \simeq H^i(M_{\leq a})$  (the case  $i = 0$  needs a separate discussion). For real (or rational) coefficients: replace  $\mathbb{Z}$  by  $\mathbb{R}$  (or  $\mathbb{Q}$ ). Then there are two cases:  $\phi = 0$  or not.

- If  $\phi = 0$ , then  $H^k(M_{\leq b}) \simeq H^k(M_{\leq a}) \oplus \mathbb{R}$ , and the remaining gradations are not changed.
- If  $\phi \neq 0$ , then  $H^{k-1}(M_{\leq b}) \simeq \ker(\phi)$ , and the remaining gradations are not changed.

4.6 Corollary: If all the cells are of even dimension, then

$$H^{odd}(M) = 0, \quad H^{2k}(M) \simeq \mathbb{Z}^{\# \text{ of } 2k \text{ cells}}.$$

4.7 Suppose  $M \subset \mathbb{R}^N$  is a compact submanifold, let  $f_q(x) = \text{dist}(q, x)^2$  for a fixed  $q \in \mathbb{R}^N \setminus M$ .

4.8 For almost all  $q \in \mathbb{R}^N$  the function  $f_q$  is Morse.

4.9 Assume that  $q = 0$ ,  $p = (a, 0, \dots, 0)$  with  $a \in \mathbb{R}_+$ ,  $T_p M = \{x_{n+1} = x_{n+2} = \dots = 0\}$ ; then  $M$  locally is the graph of a function  $g = (a + g_1, g_2, \dots, g_{N-n} : \mathbb{R}^n \rightarrow \mathbb{R}^{N-n}$ ,  $g_1(0) = 0$ ,  $g_k(0) = 0$  for  $k > 1$ ,  $Dg(0) = 0$ ;

Parametrization of  $M$ :

$$\underline{x} = (x_1, x_2, \dots, x_n) \mapsto (a + g_1(\underline{x}), g_2(\underline{x}), \dots, g_{N-n}(\underline{x}), x_1, x_2, \dots, x_n).$$

- then

$$f_q(x) = (a + g_1(\underline{x}))^2 + \sum_{j=2}^{N-n} g_j(\underline{x})^2 + \sum_{i=1}^n x_i^2 = \sum_{i=1}^n x_i^2 + 2aQ(\underline{x}) + \mathcal{O}(\|\underline{x}\|^4),$$

where  $Q$  is a quadratic form of  $g_1$ , hence

$$D^2 f_q(p) = 2(I + 2aQ).$$

Therefore

$$ind(p) = \#\{\lambda \in \text{spec } Q \mid \lambda < -\frac{1}{2a}\}$$

## Weak Lefschetz

**4.10** Lemma: If  $M \subset \mathbb{C}^N$  is a complex submanifold,  $q \notin M$  and  $p$  is a critical point of  $f_q$ , then

$$\text{index}(p) \leq \dim_{\mathbb{C}}(M)$$

- Proof: we assume as before that  $q = 0$ ,  $p = (a, 0, \dots, 0)$ ,  $a \in \mathbb{R}_+ \subset \mathbb{C}$ .
- Very easy algebraic lemma: Suppose  $Q$  is a nondegenerate quadratic form on  $\mathbb{C}^n$ . If  $v$  is an eigenvector of the real part  $\text{Re}(Q)$  with the eigenvalue  $\lambda$ , then  $iv$  is an eigenvector with the eigenvalue  $-\lambda$ . Hence the eigenvalues are symmetrically distributed with respect to 0.
- Corollary the index of  $2(I + a \text{Re}(D^2g(0))) = 2(I + 2a \text{Re}(Q))$  is at most  $\frac{1}{2} \dim_{\mathbb{R}}(M)$ .

**4.11** If  $M \subset \mathbb{C}^N$  is a complex submanifold of the complex dimension  $n$ , then  $M$  has the homotopy type of  $n$ -dimensional CW-complex. Hence  $H^k(M; R) = 0$  for  $k > n$  (with coefficient in any ring  $R$ ).

**4.12 „Weak Lefschetz” aka „Lefschetz hyperplane theorem”** [Milnor, Morse Theory §7]: If  $X \subset \mathbb{P}^N$  is a complex submanifold of dimension  $n$ ,  $i : Y = X \cap \mathbb{P}^{N-1} \rightarrow X$ , then  $X$  is a sum of  $Y$  with cells of dimension  $k \geq n$ . Thus

- $i^* : H^k(X) \rightarrow H^k(Y)$  is an isomorphism for  $k < n - 1$  and mono for  $k = n - 1$ ,
- $i_* : H_k(Y) \rightarrow H_k(X)$  is an isomorphism for  $k < n - 1$  and epi for  $k = n - 1$ .
- Moreover  $i_* : \pi_1(Y) \rightarrow \pi_1(X)$  is an isomorphism if  $2 < n$ , epimorphism if  $2 = n$ .

**4.13** If  $X \subset \mathbb{P}^N$ , and  $M$  is a smooth hypersurface of degree  $d$ , then  $M \cap X \simeq \iota(X) \cap H$ , where  $\iota : \mathbb{P}^N \rightarrow \mathbb{P}(\text{Sym}^d(\mathbb{C}^{N+1}))$  is the Veronese embedding and  $H$  is a linear hypersurface in  $\mathbb{P}(\text{Sym}^d(\mathbb{C}^{N+1}))$ .

- Hence for complete intersection  $X \subset \mathbb{P}^N$  we have information about all Betti numbers, except the middle one:

$$X = X_{N-n} \subset X_{N-n-1} \subset \dots \subset X_{N-1} \subset X_N = \mathbb{P}^N$$

$\dim(X_i) = N - i$ , since  $k < n < \dim(X_i)$  for  $i < N - n$ , we have isomorphisms  $H^k(X_i) \simeq H^k(x_{i+1})$ .

$$H^k(X) = \begin{cases} \mathbb{Z} & \text{for } k \text{ even} \\ 0 & \text{for } k \text{ odd} \end{cases}$$

for  $k < n$ , and from Poincaré duality  $H^k(X) \simeq H_{2n-k}(X)$  we get the same result for  $n > k$ .

**4.14** Exercise: compute  $\dim(H^n(Q_n))$  for a nonsingular quadric  $Q_n \subset \mathbb{P}^{n+1}$ .

## 5 Hodge theory

### Differential forms and de Rham cohomology – summary

**5.1** Global differential forms on a  $C^\infty$ -manifold  $M$  will be denoted by  $A^\bullet(M) = \bigoplus_{k=0}^{\dim M} A^k(M)$ . (The notation  $\Omega^\bullet(M)$  is reserved for holomorphic forms.)

**5.2**  $A^\bullet(M)$  is a commutative algebra with gradation  $ab = (-1)^{\deg(a)\deg(b)}ba$

**5.3** differential satisfies the Leibniz rule  $d(ab) = ad(b) + (-1)^{\deg(a)}b$

**5.4** the linear space  $A^k(M)$  is the space of the global sections of a sheaf  $A_M^k$ .

**5.5**  $\mathbb{R}_M \hookrightarrow A_M^0 \rightarrow A_M^1 \rightarrow A_M^2 \rightarrow \dots$  is a soft (in particular acyclic) resolution of the constant sheaf  $\mathbb{R}_M$ , therefore

$$H^k(A^\bullet(M), d) = H^k(M; \mathbb{R}_M) \simeq H_{\text{sing}}^k(M; \mathbb{R}).$$

The cohomology groups are denoted by  $H^k(M)$ , we skip  $\mathbb{R}$  in the notation.

**5.6** exterior product of forms induces multiplication in cohomology  $H^k(M) \times H^\ell(M) \rightarrow H^{k+\ell}(M)$

**5.7** if  $M$  is compact,  $n = \dim M$  and  $M$  has a chosen orientation, then the integral of  $n$ -forms induces a map  $\int_M : H^n(M) \rightarrow \mathbb{R}$ . If  $M$  is connected, then  $\int_M$  is an isomorphism.

**5.8** (Poincaré Duality) if  $M$  is compact, oriented of dimension  $n$ , then the bilinear form

$$\int_M - \wedge - : H^k(M) \times H^{n-k}(M) \rightarrow \mathbb{R}$$

is nondegenerate.

**5.9** if  $M$  is oriented (not necessarily compact), then we consider cohomology with compact supports

$$H_c^k(M) = H^k(A_c^\bullet(M)).$$

Then

$$\int_M - \wedge - : H_c^k(M) \times H^{n-k}(M) \rightarrow \mathbb{R}$$

is defined and it is a nondegenerate 2-linear form.

**5.10** Having a Riemannian metric on a compact manifold allows to define **harmonic** forms  $\mathcal{H}^k(M)$  (see 5.17). The harmonic forms are closed and the resulting map  $\mathcal{H}^k(M) \rightarrow H^k(M)$  is an isomorphism. However the product of harmonic forms does not have to be harmonic.

## Hodge theory for $C^\infty$ manifolds

Suppose  $M$  is equipped with Riemannian metric, i.e. a scalar product at each tangent space  $T_x M$ . Let  $n = \dim M$ .

**5.11** Volume form is denoted by  $vol \in A^n(M)$ .

**5.12** Hodge star: for  $x \in M$

$$* : \Lambda^k T_x^* M \rightarrow \Lambda^{n-k} T_x^* M$$

It is defined by the property

$$a \wedge *b = \langle a, b \rangle vol$$

for each  $a, b \in \Lambda^k T_x^* M$ . The Hodge star extends to

$$* : A^k(M) \rightarrow A^{n-k}(M)$$

pointwise.

**5.13** We have

- (i)  $*^2 = (-1)^{k(n-k)}$  on  $k$ -forms.
- (ii)  $\langle \alpha, *\beta \rangle = (-1)^{k(n-k)} \langle *\alpha, \beta \rangle$ ,

**5.14** Let's define  $d^* = (-1)^{n(k+1)+1} * d * : A^k(M) \rightarrow A^{k-1}(M)$ .

**5.15** For compact manifold  $M$ ,  $a \in A^{k-1}(M)$ ,  $b \in A^k(M)$  we have

$$\langle da, b \rangle_M = \langle a, d^*b \rangle_M$$

We say that  $d^*$  is formally adjoint to  $d$ .

Proof

$$0 = \int_M d(a \wedge *b) = \int_M da \wedge *b + (-1)^{k-1} \int_M a \wedge d(*b).$$

Hence

$$\int_M da \wedge *b = (-1)^k \int_M a \wedge d(*b).$$



$$\begin{aligned}
\langle a, d^*b \rangle_M &= \int_M \langle a, (-1)^{d(k+1)+1} * d * b \rangle vol \\
&= (-1)^{d(k+1)+1} \int_M a \wedge * * d * b & \deg(*d * b) = k - 1 \\
&= (-1)^{d(k+1)+1+(k-1)(d-k+1)} \int_M a \wedge d * b & d(k+1) + 1 + (k-1)(d-k+1) \equiv_2 k \\
&= (-1)^k \int_M a \wedge d(*b) = \int_M da \wedge *b = \int_M \langle da, b \rangle vol
\end{aligned}$$

**5.16** Laplacian on forms is defined by

$$\Delta = dd^* + d^*d$$

It can be interpreted as the „super-commutator”  $[d, d^*]_s$ .

- In general the supercommutator of elements of a graded algebra  $A = \bigoplus_{k \in \mathbb{Z}} A^k$  is defined by

$$[\phi, \psi]_s = \phi\psi - (-1)^{k\ell}\psi\phi \quad \text{if} \quad \phi \in A^k, \quad \psi \in A^\ell.$$

**5.17** Harmonic forms:  $\mathcal{H} := \ker \Delta$ .

**5.18** The operator  $\Delta = dd^* + d^*d$  is formally self-adjoint

$$(\Delta a, b) = (a, \Delta b).$$

**5.19** For a compact oriented  $C^\infty$ -manifold  $M$  the following holds in  $A^\bullet(M)$

- 1)  $\mathcal{H} = \ker(d) \cap \ker(d^*)$
- 2)  $\ker(d^*) = \text{im}(d)^\perp$ ,  $\ker(d) = \text{im}(d^*)^\perp$ ,  $\ker(\Delta) = \text{im}(\Delta)^\perp$ ,  
(hence  $\mathcal{H} = \ker(d) \cap \text{im}(d)^\perp$ )
- 3) the spaces  $\mathcal{H}$ ,  $\text{im}(d)$  and  $\text{im}(d^*)$  are perpendicular.

Proof:

- 1) suppose  $a \in \ker(\Delta)$ :

$$0 = (\Delta a, a) = (dd^*a, a) + (d^*da, a) = (d^*a, d^*a) + (da, da) = \|d^*a\|^2 + \|da\|^2$$

- 2) Let  $P = d, d^*$  of  $\Delta$ . If  $a \in \ker(P^*)$  then  $0 = (P^*a, b) = (a, Pb)$ , hence  $a \in \text{im}(P)^\perp$ .

Conversely, if  $a \in \text{im}(P)^\perp$ , then  $0 = (a, PP^*a) = \|P^*a\|^2$ , so  $P^*a = 0$ .

- 3) It remains to show that the spaces  $\text{im}(d)$  and  $\text{im}(d^*)$  are perpendicular  $(d^*a, db) = (a, d^2b) = 0$ .  
(Here we used that  $d^2 = 0$ , all the rest was an abstract properties of formally adjoint operators.)

**5.20** Hodge decomposition

$$\mathcal{A}^\bullet(M) = \underbrace{\text{im}(d) \oplus \mathcal{H}}_{\ker(d)} \oplus \text{im}(d^*).$$

This decomposition is orthogonal.

- The decomposition follows from a general property of elliptic differential operators, which we will not prove. We would have to extend the space of  $C^\infty$  forms and consider Sobolev spaces. See [C. Voisin, Hodge Theory And Complex Algebraic Geometry I, Cambridge Studies in Advanced Mathematics. Theorem 5.22, p.128-9]. For any *elliptic* operator  $P : C^\infty(E) \rightarrow C^\infty(F)$

$$C^\infty(E) = \ker(P) \oplus P^*(C^\infty(F)).$$

(Exercise: prove the corresponding statement for a linear map between finite dimensional spaces.)

- In our case  $P = \Delta$ ,  $P^* = \Delta$

$$\mathcal{A}^\bullet(M) = \mathcal{H} \oplus \Delta(\mathcal{A}^\bullet(M)).$$

Moreover we have

$$\text{im}(\Delta) \subset \text{im}(d) + \text{im}(d^*).$$

But from orthogonality  $(\text{im}(d) \oplus \text{im}(d^*)) \cap \mathcal{H} = 0$ , hence

$$\text{im}(\Delta) = \text{im}(d) \oplus \text{im}(d^*).$$

**5.21** Corollary 1:  $\mathcal{H} \rightarrow H^*(M)$  is an isomorphism.

Moreover: if  $\Delta(a) = 0$  and  $a' = a + db$ , then  $\|a'\| \geq \|a\|$ .

Any harmonic form is the representative of its cohomology class, which has the smallest norm.

**5.22** Corollary 2: Tricky proof of the Poincaré duality: Let  $[\alpha] \neq 0 \in H^*(M)$ , then there exists a class  $[\beta]$  (in the complementary gradation) such that  $\int_M \alpha \wedge \beta \neq 0$ .

• Proof: let's assume that  $\alpha$  is harmonic. Set  $\beta = *\alpha$ . Then  $\beta$  is harmonic as well ( $d(*\alpha) = \pm *d^*(\alpha) = 0$  and  $d^*(\alpha) = \pm *d(\alpha) = 0$ ). We have

$$\int_M \alpha \wedge *\alpha = \int_M \|\alpha\|^2 \text{vol} = \|\alpha\|_M^2.$$

**5.23** Heat equation  $\alpha : \mathbb{R}_+ \rightarrow A^*(M)$  with the initial condition  $\alpha(0) = \alpha$

$$\frac{d}{dt}\alpha(t) = -\Delta\alpha(t),$$

see [D. Arapura, Algebraic Geometry over the Complex Numbers] §8

- the solution exists for  $t \geq 0$
  - $\alpha_H := \lim_{t \rightarrow \infty} \alpha(t)$  exists and is a harmonic form.
- (Laplacian has nonnegative eigenvalues: if  $\Delta(\alpha) = \lambda\alpha$  then

$$\lambda\|\alpha\| = (\Delta\alpha, \alpha) = \|d\alpha\|^2 + \|d^*\alpha\|^2 \geq 0.$$

hence the limit exists.)

- $\alpha = \alpha_H + \Delta G(\alpha)$ , where  $G(\alpha) = \int_0^\infty (\alpha(t) - \alpha_H) dt$  is the Green operator  $G : \mathcal{H}^\perp \rightarrow A^\bullet(M)$ .
- Let's check for  $\alpha$  being an eigenvector  $\Delta\alpha = \lambda\alpha$ ,  $\lambda \neq 0$ : The solution is of the form  $\alpha(t) = e^{-\lambda t}\alpha$ . Then

$$\Delta \left( \int_0^\infty e^{-\lambda t} \alpha dt \right) = \int_0^\infty e^{-\lambda t} \lambda \alpha dt = \left( \int_0^\infty e^{-\lambda t} \lambda dt \right) \alpha = \alpha.$$

- If  $\beta(t)$  is a solution with the initial condition  $\beta$ , then  $d\beta(t)$  is a solution with the initial condition  $d\beta$  (because  $d\Delta = ddd^* + dd^*d = dd^*d + dd^*d + d^*dd = \Delta d$ ).
  - If  $\alpha = \alpha_H + d\beta$  then  $\alpha_t = \alpha_h + d\beta_t$ .
  - If  $d\alpha = 0$ , then  $d\alpha_t = 0$  and  $[\alpha_t] = [\alpha]$
- Proof  $\alpha = \alpha_h + d\beta$ ,  $(\alpha_t - \alpha_h)' = -\Delta(d\beta_t) = -d\Delta(\beta_t)$

## Hermitian linear algebra

Suppose  $(V, I)$  is a real vector space with a complex structure.

**5.24** Hermitian product

$$V \otimes V \rightarrow \mathbb{C}$$

$$\langle\langle v, w \rangle\rangle = \langle v, w \rangle - i\omega(v, w)$$

consists of:

- $I$ -invariant scalar product  $\langle v, w \rangle$ ,
- $I$ -symplectic form  $\omega(v, w)$
- the scalar product and the symplectic form determine each other  $\omega(v, w) = \langle I(v), w \rangle = -\langle v, I(w) \rangle$ .

**5.25** The volume form is defined as the wedge of an orthonormal (positively oriented) basis vectors of  $V^*$ :

- suppose  $\dim_{\mathbb{C}}(V) = n$

$$\text{vol} = (dx_1 \wedge dy_1) \wedge \cdots \wedge (dx_n \wedge dy_n) = \left(\frac{i}{2}\right)^n (dz_1 \wedge d\bar{z}_1) \wedge \cdots \wedge (dz_n \wedge d\bar{z}_n)$$

•

$$\omega = \sum_{k=1}^n dx_k \wedge dy_k = \frac{i}{2} \sum_{k=1}^n dz_k \wedge d\bar{z}_k.$$

$$\omega^n = n! \text{vol}.$$

**5.26**  $\omega$  as a differential form on  $\mathbb{C}^n$  is closed and  $U_n$  invariant.

## 6

**6.1** General picture:

- 1) Manifolds with Riemannian metric  $\leadsto$  harmonic forms represent cohomology classes
- 2) Complex manifolds  $\leadsto$  complex coordinates, forms  $dz$  and  $d\bar{z}$ , decomposition of differential forms into types  $(p, q)$
- 1) & 2) hermitian manifolds  $\leadsto$  the differential form  $\omega$  of type  $(1,1)$
- 3) Kähler manifolds (the condition  $d\omega = 0$ )  $\leadsto$  decomposition of cohomology into types and  $\mathfrak{sl}_2$  action.

### Main example - the projective space

**6.2** The projective space  $\mathbb{P}^n$  can be obtained as the quotient  $S^{2n+1}/S^1$ . The tangents space  $T_{[z]}\mathbb{P}^n = T_z S^{2n+1}/T_z(S^1 z)$ .

**6.3** the form  $\omega$  is well defined on the quotient space: tangent vector space  $T_z(S^1 z)$  is spanned by the vector  $(\frac{d}{dt}e^{it}z)_{t=0} = Iz$ . Therefore for  $w \in T_z S^{2n+1} = z^\perp$

$$\omega(w, Iz) = \langle Iw, Iz \rangle = \langle w, z \rangle = 0.$$

Since  $\omega$  is  $S^1$  invariant the choice of  $z \in [z]$  leads to the same form.

**6.4** We define a 2-form  $\omega_{FS}(w_1, w_2) = \omega(\tilde{w}_1, \tilde{w}_2)$ , where  $\tilde{w}_1, \tilde{w}_2$  are any lifts of  $w_1, w_2 \in T_{[z]}\mathbb{P}^n$  to  $T_z S^{2n+1}$ . Let  $p : S^{2n+1} \rightarrow \mathbb{P}^n$  be the projection.

- The form  $\omega_{FS}$  satisfies  $p^*(\omega_{FS}) = \omega$ .
- The form  $\omega_{FS}$  is closed because  $p^*$  is injective on forms and  $d\omega = 0$ .

**6.5** For  $n = 1$ , on  $U_0 = \{z_0 \neq 0\} \simeq \mathbb{C}$  there is a section

$$(s_1, s_2) : U_0 \rightarrow S^3 \subset \mathbb{C}^2$$

$$s_1(z) = \frac{1}{\sqrt{1+|z|^2}}, \quad s_2(z) = \frac{z}{\sqrt{1+|z|^2}}.$$

Then

$$\omega_{FS}(z) = \frac{i}{2}(s_1^*(dz_1 \wedge d\bar{z}_1) + s_2^*(dz_2 \wedge d\bar{z}_2))$$

Since the image of  $s_1$  is contained in  $\mathbb{R}$ , thus  $s_1^*(dz_1 \wedge d\bar{z}_1) = 0$ . The second summand is equal to Jacobian times  $dx \wedge dy$ ,

$$(x, y) \mapsto \left( \frac{x}{\sqrt{1+x^2+y^2}}, \frac{y}{\sqrt{1+x^2+y^2}} \right)$$

$$J(x, y) = \det \left( \frac{1}{(1+x^2+y^2)^{3/2}} \begin{bmatrix} 1+y^2 & -xy \\ -xy & 1+x^2 \end{bmatrix} \right)$$

Hence

$$\omega_{FS}(z) = \frac{1}{(1+x^2+y^2)^2} dx \wedge dy.$$

The volume of  $\mathbb{P}^1$ :

$$\int_{\mathbb{R}^2} \frac{1}{(1+x^2+y^2)^2} dx \wedge dy = 2\pi \int_{\mathbb{R}_+} \frac{r}{(1+r^2)^2} dr \stackrel{r^2=u}{=} \pi \int_{\mathbb{R}_+} \frac{1}{(1+u)^2} du = \pi.$$

**6.6** Often we normalize

$$\omega_{FS} := \frac{1}{\pi} \omega_{FS}.$$

With this normalization  $\int_{\mathbb{P}^n} \omega_{FS}^n = 1$ .

**6.7** The class  $[\omega_{FS}] \in H^2(\mathbb{P}^n)$  is a generator of  $H^*(\mathbb{P}^n; \mathbb{R}) \simeq \mathbb{R}[h]/(h^{n+1})$

**6.8** The normalized class  $[\omega_{FS}]$  is represented by the (Poincaré dual) of  $[\mathbb{P}^{n-1}]$  with  $\mathbb{P}^{n-1}$  embedded as a linear hypersurface.

## $1\frac{1}{2}$ -linear algebra

**6.9** Complex structure on a real vector space is an automorphism  $I$  satisfying  $I^2 = -id$ . It decomposes  $V_{\mathbb{C}} := V \otimes \mathbb{C}$  into eigenspaces

$$V_{\mathbb{C}} = V_i + V_{-i}.$$

Necessarily  $\dim V$  is even and one can find a real basis  $e_1, e_2, \dots, e_n, f_1, f_2, \dots, f_n$  of  $V$ , such that  $I(e_k) = f_k, I(f_k) = -e_k$ .

- The vectors  $e_k - if_k$  form a basis of  $V_i$ :  $I(e_k - if_k) = f_k + ie_k = i(e_k - if_k)$
- The vectors  $e_k + if_k$  form a basis of  $V_{-i}$ :  $I(e_k + if_k) = f_k - ie_k = -i(e_k + if_k)$

**6.10** We are more concerned about the dual space:  $\mathbb{C}$ -linear form are said to have the type (1,0)

$$\Lambda^{10}V^* := \{ \phi \in Hom_{\mathbb{R}}(V, \mathbb{C}) \mid \phi(Iv) = i\phi(v) \},$$

the antilinear forms are said to have the type (0,1)

$$\Lambda^{01}V^* := \{ \phi \in Hom_{\mathbb{R}}(V, \mathbb{C}) \mid \phi(Iv) = -i\phi(v) \},$$

We have

$$V^* \otimes \mathbb{C} = \Lambda^{10}V^* \oplus \Lambda^{01}V^*.$$

**6.11** The dual basis is denoted by

$$dx_k := e_k^*, \quad dy_k := f_k^*.$$

We define

$$dz_k := dx_k + idy_k, \quad d\bar{z}_k := dx_k - idy_k.$$

The 1-forms  $dz_k$  are the basis of  $\Lambda^{10}V^*$ , and  $d\bar{z}_k$ 's are the basis of  $\Lambda^{01}V^*$ .

**6.12** We have a  $\mathbb{C}$ -linear isomorphism  $(V^*, I) \xrightarrow{\Phi} (\Lambda^{10}V^*, i)$ ,  $\Phi(f)(v) = f(v) - if(Iv)$   
 $\Phi(If)(v) = f(Iv) - if(I^2v) = f(Iv) + if(v) = i(f(v) - if(Iv)) = i\Phi(f)(v)$

And an anti-linear isomorphism:  $(V^*, I) \xrightarrow{\Psi} (\Lambda^{01}V^*, i)$ ,  $\Psi(f)(v) = f(v) + if(Iv)$   
 $\Psi(If)(v) = f(Iv) + if(I^2v) = f(Iv) - if(v) = -i(f(v) + if(Iv)) = -i\Psi(f)(v)$

**6.13** The exterior forms of the type  $(p, q)$ :

$$\Lambda^k V_{\mathbb{C}}^* = \bigoplus_{p+q=k} \Lambda^{pq}, \quad \Lambda^{pq} := \Lambda^p(\Lambda^{10}V^*) \wedge \Lambda^q(\Lambda^{01}V^*).$$

– Conjugation acts on  $\Lambda^k(V^* \otimes \mathbb{C}) = (\Lambda^k V^*) \otimes \mathbb{C}$ . We have

$$\overline{\Lambda^{pq}} = \Lambda^{qp}.$$

– The operator  $I$  acts on  $\Lambda^{p,q}V^*$  via multiplication by  $i^{(p-q)}$

**6.14** Remark: the form  $\omega$  belongs to  $\Lambda^2 V^* \cap \Lambda^{11} V^* \subset \Lambda^2 V_{\mathbb{C}}^*$ .

**6.15** Exercise  $\Lambda^{10} \perp \Lambda^{01}$

## Linear algebra on the tangent space

**6.16** Assume that  $M$  is a complex manifold, then tangent space  $T_p M$  at each point  $p \in M$  is a complex vector space. We treat it as a real vector space with an automorphism  $I$  given by the multiplication by  $i$ . Globally  $I \in \text{End}(TM)$ , i.e. is an endomorphism of the tangent bundle.

**6.17** Our method: Linear algebra  $\rightsquigarrow$  differential/complex manifolds structure

**6.18** An almost complex manifold  $(M, I)$  is a pair, where  $M$  is a real  $C^\infty$ -manifold and  $I \in \text{End}(TM)$  a tensor satisfying  $I^2 = -id$  (i.e. a complex structure in each  $T_p M$  smoothly depending on the point  $p \in M$ .)

[W tym roku nie będziemy rozważać rozmaitości niemal zespolonych w ogólności, ale od razu zakładamy, że mamy rozmaitość zespoloną. Patrz [Huybrechts §1.2], w szczególności [Huybrechts 2.6.19]]

**6.19** The eigenspace of  $I$  acting on  $T^*M \otimes \mathbb{C}$  decomposes this bundle into a direct sum of complex subbundles:

$$(T^*M \otimes \mathbb{C})_i \oplus (T^*M \otimes \mathbb{C})_{-i}.$$

- The global sections of the above bundles will be denoted by  $A^{10}(M)$  and  $A^{01}(M)$ .
- Locally a form in  $A^{10}(M)$  can be written as  $\sum_k a_k(z) dz_k$ . If we change the coordinate chart it can be written in the same form. This is because for a holomorphic map  $\phi : U' \rightarrow U$  the composition  $z_k \circ \phi : U' \rightarrow \mathbb{C}$  is holomorphic, so  $d(z_k \circ \phi) = \sum_k a'_k(z') dz'_k$  for some functions  $a'_k(z')$ .

**6.20** The complexified space of forms decomposes as a direct sum  $A^k(M)_{\mathbb{C}} = \bigoplus_{p+q=k} A^{p,q}(M)$ .

- $(p, q)$ -form locally can be written as

$$\sum_{|A|=p} \sum_{|B|=q} a_{A,B}(z) dz_A \wedge d\bar{z}_B$$

## Hermitian structure

Assume that  $M$  has a hermitian structure, this is equivalent of having Riemannian metric, which is  $I$ -invariant.

**6.21** The form  $\omega$  is of the type  $(1,1)$ , in addition it has real coefficients.

**6.22** The Lefschetz operator

$$L(\alpha) := \omega \wedge \alpha.$$

**6.23** Suppose  $\dim V = n$ . Let us define  $H \in \text{End}(\Lambda^k V)$  as the multiplication by  $k - n$  on  $\Lambda^k V$ .

- We have  $[H, L] = 2L$ .

**6.24** Let us define the adjoint operator  $L^*$

$$\langle L\alpha, \beta \rangle = \langle \alpha, L^*\beta \rangle$$

lowering the gradation by 2. We have:

- $[H, L] = 2L, \quad [H, L^*] = -2L^*$
- $[L, L^*] = H$ .

**6.25** The vector space of forms at a point  $p \in M$ , i.e.  $\bigoplus_k \Lambda^k T_p^*(M)$  is a representation of the Lie algebra  $\mathfrak{sl}_2(\mathbb{Z})$ . We obtain a representation on the global forms.

$$\rho : \mathfrak{sl}_2(\mathbb{Z}) \rightarrow \text{End}(A^\bullet(M)), \quad \rho(h) = H, \quad \rho(\ell) = L, \quad \rho(\ell^*) = L^*,$$

where

$$h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \ell = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \ell^* = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

## Recollection from representation theory

See e.g. [Fulton-Harris, Representation Theory: A First Course, §11]

- Representation of a Lie algebra  $\mathfrak{g}$  on  $V$  is (by definition) a morphism of Lie algebras  $\mathfrak{g} \rightarrow \text{End}(V)$ .
- Having a representation of  $\mathfrak{sl}_2(\mathbb{Z})$  is equivalent to having three linear maps  $L, H, L^*$  such that

$$[H, L] = 2L, \quad [H, L^*] = -2L^* \quad [L, L^*] = H.$$

It costs nothing to extend linearly such representation to  $\mathfrak{sl}_2(K)$  if  $V$  is a vector space over the field  $K = \mathbb{R}$  or  $\mathbb{C}$ .

- Any finite dimensional representation of  $\mathfrak{sl}_2(\mathbb{Z})$  is a direct sum of simple subrepresentations. („Simple” means that it has no nontrivial subrepresentations.)
- Simple representations are of the form  $S_k = \text{Sym}^k(\mathbb{C}^2)$  (the same for the theory over  $\mathbb{R}$ ). Other description:

$$S_k = \{\text{homogeneous polynomials of degree } k \text{ in variables } x, y\}$$

$$\ell(f) = y \frac{df}{dx}, \quad \ell^*(f) = x \frac{df}{dy}.$$

**6.26** With the assumption that  $d\omega = 0$  we will show that  $\mathfrak{sl}_2$  action on forms induces an action on cohomology and deduce very important consequences.

## 7 Differential on complex manifolds

**7.1** If  $M$  is a complex manifold, then

$$d(A^{p,q}(M)) \subset A^{p+1,q}(M) \oplus A^{p,q+1}(M)$$

$$d = \partial + \bar{\partial}, \quad \partial^2 = 0 = \bar{\partial}^2, \quad \partial\bar{\partial} + \bar{\partial}\partial = 0.$$

**7.2** Let  $\Omega^p(M)$  denote the form of the type  $(p, 0)$  with holomorphic coefficients.

- Lemma:

$$\Omega^p(M) = \ker(\bar{\partial} : A^{p,0}(M) \rightarrow A^{p,1}(M)).$$

**7.3** Dolbeault complex: for  $0 \leq p \leq \dim_{\mathbb{C}} M$  we have a complex

$$0 \rightarrow A^{p,0}(M) \xrightarrow{\bar{\partial}} A^{p,1}(M) \xrightarrow{\bar{\partial}} \dots \xrightarrow{\bar{\partial}} A^{p,\dim M}(M) \rightarrow 0,$$

**7.4** We define Dolbeault cohomology [Huybrechts 2.6.20]:

$$H_{\text{Dol}}^q(M; \Omega^p) := H^q(A^{p,\bullet}(M), \bar{\partial})$$

**7.5** Holomorphic Poincaré lemma [Huybrechts 1.3.7]: the complex of sheaves on  $M$

$$0 \rightarrow \Omega^p \rightarrow A^{p,0} \rightarrow A^{p,1} \rightarrow A^{p,2} \rightarrow \dots$$

is exact.

- This means that if  $\bar{\partial}\alpha = 0$ ,  $\alpha \in A^{p,q}(U)$ , then *locally* there exists  $\beta$  such that  $\bar{\partial}\beta = \alpha$ , i.e. for each point  $p \in U$  there exists  $V \subset U$ ,  $p \in V$  and  $\beta \in A^{p,q-1}(V)$  such that  $\bar{\partial}\beta = \alpha|_V$ .

**7.6** It is enough to solve the following problem:

- Holomorphic Poincaré lemma in 1 variable: Let  $\mathbb{D}_{\varepsilon} \subset U \subset \mathbb{C}$ , where  $U$  is open, and let  $f \in C^{\infty}(U; \mathbb{C})$  be a smooth function. Suppose  $\frac{\partial}{\partial \bar{z}} f = 0$ , then there exists  $g \in C^{\infty}(\mathbb{D}_{\varepsilon}; \mathbb{C})$  such that  $\frac{\partial}{\partial \bar{z}} g = f$ .
- The solution to the previous problem, with  $f = f_w$  depending smoothly on a parameter can be found in a way that  $g_w$  depends smoothly on the parameter.

$$g(z) = \mathcal{I}(f)(z) = \frac{1}{2\pi i} \int_{\mathbb{D}_\varepsilon} \frac{f(\xi)}{\xi - z} d\xi \wedge d\bar{\xi}.$$

- Analogy with the real case:

— for a real (compactly supported)  $f : \mathbb{R} \rightarrow \mathbb{R}$  we define the primitive function

$$\mathcal{I}(f)(x) = \int_{-\infty}^x f(\xi) d\xi = \int_{-\infty}^{+\infty} K(\xi - x) f(\xi) d\xi,$$

where

$$K(\xi) = \begin{cases} 0 & \text{if } \xi < 0 \\ 1 & \text{if } \xi \geq 0 \end{cases} \quad \text{and} \quad K'(\xi) = \delta_0.$$

So the primitive function is expressed by the convolution with  $K$ , i.e  $\mathcal{I}(f)(x) = (K * f)(x)$ .

(In general  $(f_1 * f_2)' = f_1' * f_2$ .)

— similarly for complex, compactly supported function  $f : \mathbb{C} \rightarrow \mathbb{C}$

$$g(z) = (K * f)(z),$$

where  $K(z) = \frac{1}{2\pi i} \frac{1}{z}$ , which has the property  $\frac{\partial}{\partial \bar{z}} K = \delta_0$

## Sheaf cohomology - a summary, see eg [Huybrechts, Appendix B]

**7.7** Cohomology with the coefficients in a sheaf  $\mathcal{F}$ : there are two important construction

- Čech cohomology
  - Sheaf cohomology as the derived functor of  $\Gamma$  - taking the global sections.
- 1) we find a resolution of  $\mathcal{F}$ , i.e. an exact complex

$$0 \rightarrow \mathcal{F} \rightarrow I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow \dots$$

with the sheave  $A^k$  sufficiently good (acyclic, e.g. injective)

2) we apply the functor of global sections (and cut off the first term)

$$\Gamma(I^0) \rightarrow \Gamma(I^1) \rightarrow \Gamma(I^2) \rightarrow \dots$$

This complex is no longer exact.

3) We compute cohomology:

$$H^k(M; \mathcal{F}) = H^k(\Gamma(I^\bullet)).$$

We have  $H^0(M; \mathcal{F}) = \Gamma(\mathcal{F})$ , because the functor  $\Gamma$  is left-exact.

**7.8** In our case, when the base is paracompact any *soft* resolution is acyclic. („Soft” means, that sections defined on a closed set can be extended to global sections.)

- Suppose  $M$  is a  $C^\infty$ -manifold. Any sheaf which is a module over the ring of  $C^\infty$ -functions is soft.
- The complex of  $C^\infty$ -forms on  $C^\infty$ -manifold  $A^0 \rightarrow A^1 \rightarrow A^2 \rightarrow \dots$  is a resolution of the sheaf  $\ker(d : A^0 \rightarrow A^1) = \underline{\mathbb{R}}_M$ , the sheaf of locally constant functions.

**7.9** The sheaves  $A^{p,q}$  are  $A^0$ -modules, hence they are soft.

- The Dolbeault complex is a resolution of  $\Omega_M^p = \ker(\bar{\partial} : A^{p,0} \rightarrow A^{p,1})$
- 

$$H^k(M; \Omega^p) = H^k(A^{p,\bullet}(M))$$

i.e the Dolbeault cohomology is the sheaf cohomology in the sense of the homological algebra.

**7.10** If  $M$  is a complex manifold, then  $A_{\mathbb{C}}^{\bullet} = \bigoplus_{p+q=\bullet} A^{p,q}$  is a resolution of the sheaf  $\mathbb{C}_M$ .

- For  $p \geq 0$  define the Hodge's filtration (on the sheaf level)

$$F^p A^k = \bigoplus_{p'+q=k, p' \geq p} A^{p',q}.$$

Claim:  $F^p A^{\bullet}$  is a subcomplex of  $A^{\bullet}$ .

- The resulting filtration in cohomology  $H^k(M; \mathbb{C}) = H^k(A^{\bullet}(M)_{\mathbb{C}})$

$$F^p H^k(M; \mathbb{C}) = \text{im}(H^k(F^p A^{\bullet}(M)) \rightarrow H^k(A^{\bullet}(M)_{\mathbb{C}})).$$

**7.11** We have

$$F^p A^k / F^{p+1} A^k \simeq A^{p,k-p}.$$

- The quotient map is a map of complexes (with a shift of the gradation)

$$(F^p A^{p+\bullet}, d) \rightarrow (A^{p,\bullet}, \bar{\partial})$$

- We have maps of complexes (I denote the shift of gradations by  $[i]$ . i.e.  $(F[i]^k = F^{k+i})$

$$A^{\bullet} \leftarrow F^p A^{\bullet} \rightarrow A^{p,\bullet}[-p]$$

- Passing to cohomology:

$$H^k(M; \mathbb{C}) \leftarrow H^k(M; F^p A^{\bullet}) \rightarrow H^{k-p}(X; \Omega^p).$$

**7.12** The relation between cohomologies of the quotients with cohomology of the entire sheaf is given by the spectral sequence

$$E_1^{p,q} = H^{p+q}(F^p A^{\bullet}(M) / F^{p+1} A^{\bullet}(M)) = H^q(M; \Omega_M^p) \Rightarrow H^{p+q}(M; \mathbb{C}).$$

## Generalities about spectral sequence

If  $C^{\bullet}$  is a complex with decreasing filtration

$$C^{\bullet} = F^0 C^{\bullet} \supset F^1 C^{\bullet} \supset F^2 C^{\bullet} \supset \dots,$$

then one wishes to relate cohomologies  $H^*(F^p C^{\bullet} / F^{p+1} C^{\bullet})$  with  $H^*(C^{\bullet})$ .

- There exists a spectral sequence (under some boundness of degree assumptions)

$$E_0^{p,q} = F^p C^{p+q} / F^{p+1} C^{p+q}, \quad E_1^{p,q} = H^{p+q}(F^p C^{\bullet} / F^{p+1} C^{\bullet}), \quad \dots$$

- There exists a sequence of tables  $E_r^{p,q}$  with differentials of degree  $(1-r, r)$ , such that

- 1)  $H^*(E_r^{\bullet,\bullet}) = E_{r+1}^{\bullet,\bullet}$
- 2)  $E_{\infty}^{p,q} = F^p H^{p+q}(C^{\bullet}) / F^{p+1} H^{p+q}(C^{\bullet})$

**7.13** For the total complex of the bicomplex  $A^{p,q}(M)$  with the Hodge filtration  $F^p A^{\bullet}(M) = A^{\geq p,\bullet}(M)$  the resulting spectral sequence is called the **Frölicher spectral sequence**.

## Hodge theory for Hermitian manifolds

**7.14** Hermitian structure on a complex manifold  $M$  is a choice of a Hermitian product in each tangent space.

- such structure is a section of  $T^*M \otimes \bar{T}^*M$  which is symmetric and positively definite. We assume that it is a  $C^{\infty}$
- real part is a scalar product, the imaginary part - a differential 2-form (which does not have to be closed).
- Hermitian structures exist for paracompact manifolds: we can chose a Hermitian structure locally in maps and glue them using partition of unity.



**7.15** We extend Hodge  $*$   $\mathbb{C}$ -linearly

- If  $\dim M = 1$

$$*dz = *(dx+idy) = dy-idx = -i(dx+idy) = -idz, \quad *d\bar{z} = *(dx-idy) = dy+idx = i(dx-idy) = id\bar{z}$$

$$*1 = \omega = dx \wedge dy = \frac{i}{2} dz \wedge d\bar{z}, \quad *\omega = 1$$

- In higher dimensions

$$* : \Lambda^{p,q} \xrightarrow{\cong} \Lambda^{n-q,n-p}$$

$$*dz_I \wedge d\bar{z}_J = c dz_{[n] \setminus J} \wedge d\bar{z}_{[n] \setminus I}$$

Exercise: compute  $c$ .

- Occasionally will appear antilinear star

$$\bar{*} : \Lambda^{p,q} \xrightarrow{\cong} \Lambda^{n-p,n-q}, \quad \bar{*}(\alpha) = *\bar{\alpha} = \overline{* \alpha}.$$

**7.16** We have operators real  $L, L^*, H = [L, L^*] = (\deg - n)id$  acting on  $C^\infty$ -forms  $A^*(X)$ . The adjoint operator

$$L^* = *^{-1} L * = (-1)^{\deg} * L *$$

(The sign should be  $(-1)^{(\dim_{\mathbb{R}} M - \deg) \deg}$  but here  $\dim_{\mathbb{R}} TM$  is even). Often in literature  $L^*$  is denoted by  $\Lambda$ , but it can be confused with the exterior power). The adjoint operator satisfies  $(L\alpha, \beta) = (\alpha, L^*\beta)$ .

- The complexified operators  $L, L^*, H = [L, L^*] = (\deg - n)id$  act on  $A^*(X)_{\mathbb{C}}$ . Hence  $A^*(X)_{\mathbb{C}}$  becomes a (infinite dimensional) representation of  $\mathfrak{sl}(2)$ .
- We take complexification, because we are also interested in the bigradation, available only over  $\mathbb{C}$ .

**7.17** We define operators

$$\partial^* = - * \bar{\partial} * : A^{p,q}(X) \rightarrow A^{p-1,q}(X),$$

$$(p, q) \mapsto (n - q, n - p) \mapsto (n - q, n - p + 1) \mapsto (p - 1, q)$$

and

$$\bar{\partial}^* = - * \partial * : A^{p,q}(X) \rightarrow A^{p,q-1}(X).$$

We have  $d^* = \partial^* + \bar{\partial}^*$ .

- explanation of signs:  $d^* = (-1)^{\dim_{\mathbb{R}} M (\deg + 1) + 1} * d * = - * d *$

### 7.18 Kähler structure

It can be defined in equivalent ways:

- Definition 1: locally there exists local coordinates in which  $\omega = \frac{i}{2} \sum_k dz_k \wedge d\bar{z}_k + \mathcal{O}(\|x\|^2)$ .  
i.e. in some coordinates the Hermitian metric is the same as for flat the manifold  $\mathbb{C}^n$  up to the terms of order 2.
- Definition 2:  $d\omega = 0$
- Proof 1)  $\Rightarrow$  2) obvious.

**7.19** Proof 2)  $\Rightarrow$  1) [C. Voisin, Hodge Theory And Complex Algebraic Geometry I, Prop 3.14]

- How to construct good coordinates?

$$\omega = \frac{i}{2} \sum_k dz_k \wedge d\bar{z}_k + \sum_{k,l} (\varepsilon_{k,l}^h + \varepsilon_{k,l}^a) dz_k \wedge d\bar{z}_l + \mathcal{O}(|z|^2)$$

where  $\varepsilon_{k,l}^h$  is a holomorphic linear form,  $\varepsilon_{k,l}^a$  antiholomorphic liner form.

- $\overline{\varepsilon_{k,l}^a} = \varepsilon_{l,k}^h$  since  $\omega$  is real.
- $\frac{\partial}{\partial z_j} \varepsilon_{k,l}^h = \frac{\partial}{\partial z_k} \varepsilon_{j,l}^h$  since  $\omega$  is closed

### 7.20 Hodge identities:

- i)  $[\bar{\partial}, L] = [\partial, L] = 0$  (since  $\omega$  is closed)
- i') equivalently  $[L^*, \partial^*] = [L^*, \bar{\partial}^*] = 0$
- ii)  $[\bar{\partial}^*, L] = i\partial$ ,  $[\partial^*, L] = -i\bar{\partial}$
- ii') equivalently  $[L^*, \bar{\partial}] = -i\partial^*$ ,  $[L^*, \partial] = i\bar{\partial}^*$  (this is the most difficult, the rest follows)
- iii)  $[\partial, \bar{\partial}^*]_s = [\partial^*, \bar{\partial}]_s = 0$  (i.e.  $\partial\bar{\partial}^* + \bar{\partial}^*\partial = 0$  etc, this is a formal consequence of ii))
- iv)  $\Delta_\partial = \Delta_{\bar{\partial}} = \frac{1}{2}\Delta$  and it commutes with  $\partial, \bar{\partial}, \partial^*, \bar{\partial}^*, L$  i  $L^*$  (formal algebraic proof)

**7.21** Short proof from [C. Voisin, Hodge Theory And Complex Algebraic Geometry I, Prop 6.5].

- Assume according to Definition 1) that  $\omega$  has a standard form up to the terms of order 2. Therefore in calculations involving only the **first derivatives** at a point we can assume that

$$\omega = \frac{i}{2} \sum dz_k \wedge d\bar{z}_k$$

- We show ii') i.e.  $[L^*, \partial] = i\bar{\partial}^*$ . It is enough to check

$$([L^*, \partial](\alpha))_{z=0} = i(\bar{\partial}^*\alpha)_{z=0}$$

- We decompose  $\omega = \sum_k \omega_k$ ,  $\omega_k = \frac{i}{2} dz_k \wedge d\bar{z}_k$ .  
The adjoint operator  $L_k^* = (\omega_k \wedge)^*$  is expressed by the contraction of differential forms

$$L_k^* = -2i\iota_{\bar{v}_k}\iota_{v_k},$$

where  $v_k = \frac{\partial}{\partial z_k}$ ,  $\bar{v}_k = \frac{\partial}{\partial \bar{z}_k}$ .

- We decompose  $\bar{\partial} = \sum \bar{\partial}_k$ . The adjoint differentials

$$\partial_k^* = -2\frac{\partial}{\partial \bar{z}_k}\iota_{v_k}, \quad \bar{\partial}_k^* = -2\frac{\partial}{\partial z_k}\iota_{\bar{v}_k},$$

A sample of check in  $\dim=1$

$$\partial^* f dz = - * \bar{\partial}^* f dz = - * \bar{\partial}(-i f dz) = i * \frac{\partial}{\partial \bar{z}} f d\bar{z} \wedge dz = -2 \frac{\partial}{\partial \bar{z}} f * \frac{i}{2} dz \wedge d\bar{z} = -2 \frac{\partial}{\partial \bar{z}} f$$

**7.22** Second Hodge identity  $[L^*, \partial] = i\bar{\partial}^*$  for the flat metric: We decompose  $\bar{\partial} = \sum \bar{\partial}_k$  and  $L^* = \sum L_k^*$ . Show that  $\bar{\partial}_k^* = -2\iota_{\bar{v}_k}\frac{\partial}{\partial z_k}$ , where  $\bar{v}_k = \frac{\partial}{\partial \bar{z}_k}$ . Note  $\partial_\ell$  commutes with  $L_k^*$  for  $k \neq \ell$ . It remains to check  $[L_k^*, \partial_k]$  for  $\alpha = f dz_I \wedge d\bar{z}_J$ , considering 4 cases  $k \in$  or  $\notin$  to  $I$  and  $J$ . For example: suppose  $k \in I$ ,  $k \in J$ . That is  $I = \{k\} \cup I'$ ,  $J = \{k\} \cup J'$ :

$$\begin{aligned} [L_k^*, \partial_k] f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'} &= \\ L_k^* \partial_k (f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'}) - \partial_k L_k^* (f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'}) &= \\ 2i \partial_k (f dz_{I'} \wedge d\bar{z}_{J'}) &= \\ 2i \frac{\partial f}{\partial z_k} dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'} &= \\ 2i \frac{\partial f}{\partial z_k} \iota_{\bar{v}_k} (d\bar{z}_k \wedge dz_k \wedge dz_{I'} \wedge d\bar{z}_{J'}) &= \\ i \bar{\partial}^* (f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'}) &= \end{aligned}$$

- It remains to check 3 other cases.

**7.23** For a computational proof see Huybrechts.

- The Huybrechts' proof of ii'): an operator  $d^c = I^{-1}dI$  is introduced and the adjoint operator  $(d^c)^*$

$$d^c = -i(\partial - \bar{\partial}), \quad (d^c)^* = - * d^c *.$$

He shows ii')  $[L^*, d] = -(d^c)^*$ . The proof is computational, using Lefschetz decomposition into  $L^k \alpha$ , where  $\alpha$  is primitive.

## 8 Kähler identities cont.

**8.1** Proof of iii) and iv) from i)&ii)

- iii)

$$i[\partial, \bar{\partial}^*] \stackrel{ii)}{=} [\partial, [L^*, \partial]] = \partial L^* \partial - \partial^2 L^* + L^* \partial^2 - \partial L^* \partial = 0$$

- To show and iv) it is convenient to introduce the language of supercommutators  $[a, b] = ab - (-1)^{\deg(a)\deg(b)}ba$ . In that notation

$$\Delta_\partial = [\partial, \partial^*].$$

- Leibniz rule, equivalent to the graded Jacobi identity

$$[a, [b, c]] = [[a, b], c] + (-1)^{\deg(a)\deg(b)}[b, [a, c]].$$

$$[[a, b], c] = [a, [b, c]] + (-1)^{\deg(b)\deg(c)}[[a, c], b].$$

- 

$$\Delta_\partial = [\partial^*, \partial] \stackrel{ii)}{=} i[[L^*, \bar{\partial}], \partial] \stackrel{Leibniz}{=} i([L^*, \underbrace{[\bar{\partial}, \partial]}_0] - [[L^*, \partial], \bar{\partial}]) \stackrel{ii)}{=} [\bar{\partial}^*, \bar{\partial}] = \Delta_{\bar{\partial}}$$

and from iii)  $\Delta = \Delta_\partial + \Delta_{\bar{\partial}}$ .

$$[L, \Delta_\partial] = [L, [\partial, \partial^*]] \stackrel{Leibniz}{=} \underbrace{[[L, \partial], \partial^*]}_0 + [\partial, [L, \partial^*]] \stackrel{ii)}{=} i[\partial, -i\bar{\partial}] = 0$$

## Cohomology of Kähler manifold

- Corollary  $H^*(M) \simeq \mathcal{H}$  is a representation of  $\mathfrak{sl}_2(\mathbb{Z})$ .

**8.2 STRATEGY:** We obtain a list operators, decompositions etc. We have shown that this structure, initially defined on forms, survives in cohomology of a complex Kähler variety.

### Lefschetz decomposition

**8.3** Let  $W$  be a representation of  $\mathfrak{sl}_2$ ,

- the eigenspaces of  $h$  are equal  $W_{k-n} = \Lambda^k T_x^* M \otimes \mathbb{C}$ ,
- $L^k$  defines an isomorphism  $W_{-k} \rightarrow W_k$  ( $k \geq 0$ ),
- $L : W_k \rightarrow W_{k+2}$  is mono for  $k < 0$ , epi for  $k + 2 > 0$ .
- Lefschetz decomposition: For  $k \geq 0$  let us define the primitive subspace

$$P_k = \{w \in W_{-k} \mid L^{k+1}w = 0\}.$$

We have

$$W_{-k} = P_k \oplus LP_{k+2} \oplus L^2P_{k+4} \oplus \dots$$

**8.4** The primitive cohomology classes (attention at the gradation shift): for  $0 \leq k \leq n$  let us define

$$P^{n-k} = \{\alpha \in H^{n-k}(M) \mid L^{k+1}\alpha = 0\}$$

$$P^{p,q} = \mathcal{H}^{p,q} \cap P_{\mathbb{C}}^{p+q}.$$

We have

$$P_{\mathbb{C}}^{n-k} = \bigoplus_{p+q=n-k} P^{p,q}.$$

Practical consequences:

**8.5 Hard Lefschetz Theorem** Let  $M$  be a Kähler manifold of dimension  $n$  and let  $0 \leq k \leq n$ . Then

$$L^k : H^{n-k}(M) \rightarrow H^{n+k}(M)$$

is an isomorphism.

- It follows

$$\begin{aligned} \dim H^k(M) &\leq \dim H^{k+2}(M) && \text{if } k+1 \leq n, \\ \dim H^k(M) &\geq \dim H^{k+2}(M) && \text{if } k+1 \geq n. \end{aligned}$$

### 8.6 Hodge decomposition for the operator $\bar{\partial}$

$$\mathcal{A}^{p,q}(M) = \underbrace{\text{im}(\bar{\partial}) \oplus \mathcal{H}_{\bar{\partial}}^{p,q}}_{\ker(\bar{\partial})} \oplus \text{im}(\bar{\partial}^*).$$

$$\bar{\partial} : A^{p,q-1} \rightarrow A^{p,q}, \quad \bar{\partial}^* : A^{p,q+1} \rightarrow A^{p,q}.$$

- 

$$H^q(M; \Omega^p) \simeq \mathcal{H}_{\bar{\partial}}^{p,q},$$

- Since  $\Delta_{\bar{\partial}} = \frac{1}{2}\Delta$ , we have

$$\begin{aligned} \mathcal{H}_{\bar{\partial}}^{p,q} &= \mathcal{H}^{p,q}, \\ \overline{\mathcal{H}^{p,q}} &= \mathcal{H}^{q,p}, \quad * \mathcal{H}^{p,q} = \mathcal{H}^{n-q, n-p}. \end{aligned}$$

- 

$$\mathcal{H}^k = \bigoplus_{p+q=k} \mathcal{H}^{p,q}.$$

### 8.7 Hodge decomposition in cohomology

- Recall the Hodge filtration

$$F^p A^k(M) = \bigoplus_{p' \geq p, p+q=k} A^{p,q}(M)$$

and the induced filtration in cohomology

$$F^p H^k(M) = \text{im}(H^k(F^p A^\bullet(M) \rightarrow H^*(M))).$$

The definition is independent from the metric and

$$F^p H^k(M) = \text{image of } \bigoplus_{p' \geq p, p+q=k} \mathcal{H}^{p,q}.$$

- Conjugating we obtain

$$\overline{F^p H^k(M)} = \text{image of } \bigoplus_{p' \geq p, p+q=k} \mathcal{H}^{q,p}.$$

- Define

$$H^{p,q}(M) = F^p H^{p+q}(M) \cap \overline{F^q H^{p+q}(M)}.$$

This definition does not depend on the Kähler metric.

$$H^k(M) = \bigoplus_{p+q=k} H^{p,q}(M),$$

**8.8** Let  $h^{p,q} = \dim H^{p,q}(M)$ .

- Hard Lefschetz implies inequalities

$$h^{p,q} \leq \dim h^{p+1,q+1} \quad \text{if } p+q+1 \leq n,$$

$$h^{p,q} \geq \dim h^{p+1,q+1} \quad \text{if } p+q+1 \geq n,$$

- The symmetries  $h^{p,q} = h^{n-p,n-q} = h^{q,p}$  are organized in the „Hodge diamond”
- For example for  $n = 3$

$$\begin{array}{ccccccc}
 & & & h^{33} & & & \\
 & & h^{32} & & h^{23} & & \\
 & h^{31} & & h^{22} & & h^{13} & \\
 h^{30} & & h^{21} & & h^{12} & & h^{03} \\
 & h^{20} & & h^{11} & & h^{02} & \\
 & & h^{10} & & h^{01} & & \\
 & & & h^{00} & & & 
 \end{array}
 =
 \begin{array}{ccccccc}
 & & & 1 & & & \\
 & & \spadesuit & & \spadesuit & & \\
 & \diamond & & \clubsuit & & \diamond & \\
 & & \heartsuit & & \heartsuit & & \\
 \square & & \diamond & & \clubsuit & & \diamond & \square \\
 & & \diamond & & \clubsuit & & \diamond & \\
 & & \spadesuit & & \spadesuit & & \\
 & & & 1 & & & 
 \end{array}$$

- Hard Lefschetz implies inequalities

$$h^{p,q} \leq \dim h^{p+1,q+1} \quad \text{if } p+q+1 \leq n,$$

$$h^{p,q} \geq \dim h^{p+1,q+1} \quad \text{if } p+q+1 \geq n,$$

**8.9** Moreover

- If  $k = n - (p+q) \geq 0$  then  $L^k : H^{p,q}(M) \rightarrow H^{p+k,q+k}(M)$  is an isomorphism
- If  $p+q \leq n$  then

$$H^{p,q}(M) = P^{p,q}(M) \oplus L(P^{p-1,q-1}(M) \oplus L^2(P^{p-2,q-2}(M) \oplus \dots$$

- Corollary: If  $M$  Kähler and compact, then the (Frölicher) spectral sequence

$$H^q(M; \Omega^p) \Rightarrow H^{p+q}(M; \mathbb{C})$$

degenerates on  $E_1$ , i.e.

$$E_1^{p,q} = H^q(M; \Omega^p) = E_\infty^{p,q}.$$

(the higher differentials vanish).

**8.10** Corollary: Suppose  $M$  Kähler and compact: if  $\alpha \in \Omega^p(M)$  then  $\partial\alpha = 0$ .

- Holomorphic implies closed.
- This is a generalization of: global holomorphic function is constant.

**8.11** We say that  $M$  is Calabi-Yau if  $\Omega^n \simeq \mathcal{O}_M$

(according to more restrictive definitions it is assumed additionally  $H^0(M, \Omega^p) = 0$  for  $0 < p < n$ )

- Thus  $h^{n,q} = h^{0,q}$ .
- For  $n = 3$  the Hodge diamond looks like this

$$\begin{array}{ccccccc}
 & & & 1 & & & \\
 & & 0 & & 0 & & \\
 & 0 & & \clubsuit & & 0 & \\
 1 & & \heartsuit & & \heartsuit & & 1 \\
 & 0 & & \clubsuit & & 0 & \\
 & & 0 & & 0 & & \\
 & & & 1 & & & 
 \end{array}$$

- We say that  $M^*$  is a cohomological mirror of  $M$  if  $h^{p,q}(M^*) = h^{n-p,q}(M)$ .
- For 3-manifolds this means  $h^{12}(M^*) = h^{11}(M)$  i  $h^{11}(M^*) = h^{12}(M)$ .
- Problem: how to find  $M^*$ ?

### 8.12 Serre duality: the exterior product

$$\wedge : \Omega^p \times \Omega^q \rightarrow \Omega^{p+q}$$

defines a bilinear map

$$H^k(M; \Omega^p) \times H^\ell(M; \Omega^q) \rightarrow H^{k+\ell}(M; \Omega^{p+q}).$$

If  $k + \ell = p + q = n$  we obtain compose it with the integral  $\int : H^n(M; \Omega^n) \simeq H^{2n}(M; \mathbb{C}) \rightarrow \mathbb{C}$ .

- By Poincaré duality this form is nondegenerate

$$H^k(M; \Omega^p) \simeq H^{n-k}(M; \Omega^{n-p})^*.$$

- More generally: we have a nondegenerate form

$$H^k(M; E) \times H^{n-k}(M; E^* \otimes \Omega^n) \rightarrow H^n(M; \Omega^n) \rightarrow \mathbb{C}$$

for a locally free sheaf  $E$ . In particular for  $\Omega^p = E$ :

$$\Omega^{n-p} \simeq \underline{\text{Hom}}(\Omega^p, \Omega^n) = (\Omega^p)^* \otimes \Omega^n$$

and we recover the previous formula.

## 9 Signature, Cousin problems

### Signature

**9.1** If  $V$  is a real vector space with a symmetric nondegenerate form  $\phi$ , then the signature

$$\sigma(V, \phi) := \dim\{\text{maximal positive definite subspace}\} - \dim\{\text{maximal negative definite subspace}\},$$

i.e.  $\#\{+\} - \#\{-\}$  after diagonalization.

- If there exists  $Z \subset V$  such that  $Z^\perp = N$ , then  $\sigma(\phi) = 0$ .

**9.2** For oriented compact  $C^\infty$ -manifold  $M$  of dimension  $4m$  the intersection pairing in  $H^{2m}(M; \mathbb{R})$

$$[\alpha] \cdot [\beta] = \int_M \alpha \wedge \beta$$

is symmetric and nondegenerate. Its signature is called the signature of  $M$ , denoted  $\text{sgn}(M)$  or  $\sigma(M)$ .

$$\sigma(M) := \sigma(H^{2m}(M), \text{intersection form}).$$

- Instead of  $H^{2m}(M)$  we can take  $H^{\text{even}}(M)$  declaring  $\alpha \cdot \beta = 0$  if  $\deg(\alpha) + \deg(\beta) \neq \dim(M)$ .
- Exercise: the signature is multiplicative:  $\sigma(M \times N) = \sigma(M)\sigma(N)$ .
- If  $M$  is a boundary of an oriented  $4m + 1$ -manifold  $W$ , then  $\sigma(M) = 0$ .

Proof: let  $\iota : M = \partial W \rightarrow W$ . Define

$$Z = \iota^*(H^{2m}(W)) \subset \iota^*(H^{2m}(M)) = V.$$

For  $[\alpha], [\beta] \in H^{2m}(W)$  by Stokes

$$\int_M \iota^* \alpha \wedge \iota^* \beta = \int_W d(\alpha \wedge \beta) = 0.$$

It remains to show, that if

$$(*) \quad [\alpha] \cdot [\iota^* \beta] = 0 \text{ for all } [\beta] \in H^{2m}(W),$$

then  $[\alpha] = \iota^*[\tilde{\alpha}]$ .

The condition  $(*)$  is equivalent to

$$[\alpha] \in \ker(H^{2m}(M) \xrightarrow{d} H^{2m+1}(W, M) \simeq (H^{2m}(W))^*).$$

From the exact sequence

$$H^{2m}(W) \xrightarrow{\iota^*} H^{2m}(M) \xrightarrow{d} H^{2m+1}(W, M)$$

we get the conclusion.

**9.3** Instead the real intersection form we consider  $H^*(M; \mathbb{C})$  with the hermitian form. The resulting signature is the same.

**9.4** Hodge'a-Riemann relations [Huybrechts 3.3.15]: Define the hermitian form  $B(\alpha, \beta)$  on  $H^k(M)$  as:

$$B(\alpha, \beta) = \int_M \alpha \wedge \bar{\beta} \wedge \omega^{n-k}.$$

This form is symmetric or antisymmetric depending on the parity of  $k$

$$B(\alpha, \beta) = (-1)^k \overline{B(\alpha, \beta)}.$$

- It is nondegenerate: for  $\alpha \in H^k(M)$  there exists  $\beta \in H^k(M)$  such that  $B(\alpha, \beta) \neq 0$ .
- Let  $\gamma \in H^{2n-k}(M)$  such that  $\int_M \alpha \wedge \bar{\gamma} \neq 0$  (e.g.  $\gamma = \bar{*}\alpha$ )
- By Hard Lefschetz  $\gamma = L^{n-k}\beta$  for some  $\beta \in H^k(M)$

$$B(\alpha, \beta) = \int_M \alpha \wedge \bar{\gamma} \neq 0.$$

- The pairing  $B$  restricted to  $H^{p,q}(M)$  is non degenerate. The form  $\gamma = \bar{*}\alpha$  is of the type  $(n-q, n-p)$ , hence  $L^{k-n}\gamma$  is of the type  $(n-q-n+k, n-p-n+k) = (p, q)$ .

**9.5** Antisymmetric forms over  $\mathbb{C}$  can be turned into symmetric:

- if  $\phi$  jest antisymmetric, i.e.

$$\phi(a, b) = -\overline{\phi(b, a)},$$

then  $\psi(a, b) := i\phi(a, b)$  is symmetric.

- If hermitian form  $\psi$  is symmetric then  $\psi(a, a) = \overline{\psi(a, a)}$ , hence  $\psi(a, a) \in \mathbb{R}$
- We say that such form is positive definite if

$$\psi(a, a) > 0 \quad \text{for } a \neq 0.$$

**9.6 Theorem [Hodge-Riemann relations]:** Let  $k = p + q$ . The form

$$i^{p-q} \cdot (-1)^{k(k-1)/2} B(\alpha, \beta)$$

restricted to the primitive space

$$P^{p,q}(M) = P^k(M) \cap H^{p,q}(M)$$

is symmetric and positive definite.

**9.7** Proof reduces to calculations for  $\Lambda^\bullet \mathbb{C}^n$ : one has to check the sign of the form  $B_0$  restricted to  $P^{p,q} \subset \Lambda^{p,q} \subset \Lambda(\mathbb{C}^n)^* \otimes \mathbb{C}$ . Here  $B_0$  is defined by the formula

$$\alpha \wedge \bar{\beta} \wedge \omega^{n-k} = B_0(\alpha, \beta) dx_1 \wedge dy_1 \wedge \cdots \wedge dx_n \wedge dy_n.$$

- We check the following identity for  $\alpha \in P^k$ :

$$(***) \quad L^{n-k}\alpha = (-1)^{\frac{k(k-1)}{2}} (n-k)! * I(\alpha),$$

or equivalently as in [Huybrechts]

$$*L^{n-k}\alpha = (-1)^{\frac{k(k+1)}{2}} (n-k)! I(\alpha),$$

where  $I$  is the complex structure acting on  $\Lambda(\mathbb{C}^n)^* \otimes_{\mathbb{R}} \mathbb{C}$ . On the  $(p, q)$  forms it acts by the multiplication by  $i^{p-q}$ .

- We show inductively

$$L^j \alpha = (-1)^{\frac{k(k-1)}{2}} \frac{j!}{(n-k-j)!} * L^{n-k-j} I(\alpha).$$

- Having  $(***)$ :

$$\begin{aligned} \alpha \wedge \bar{\alpha} \wedge \omega^{n-k} &= \alpha \wedge L^{n-k}(\bar{\alpha}) = \alpha \wedge (-1)^{\frac{k(k-1)}{2}} (n-k)! * I(\bar{\alpha}) = \\ &= (-1)^{\frac{k(k-1)}{2}} \alpha \wedge *(n-k)! I(\bar{\alpha}) = i^{q-p} (-1)^{\frac{k(k-1)}{2}} (n-k)! \langle \alpha, \alpha \rangle \text{vol} \end{aligned}$$

**9.8 Corollary** [Huybrechts 3.3.18]: Let  $n = 2m$ . Then  $M$  is a real manifold of dimension  $4m$ . The intersection form in the middle dimension  $2m$  is symmetric. It coincides with  $B(\alpha, \beta)$ .

- The signature of  $M$  is defined as the signature of the intersection form  $H^{2m}(M)$  is equal to

$$\sum_{p+q \leq m, 2|p+q} (-1)^{\frac{k(k-1)+p-q}{2}} \dim(P^{p,q}(M))$$

- We have equality  $\dim P^{p,q}(M) = h^{p,q} - h^{p-1,q-1}$ . Using symmetries of Hodge diamond  $h^{p,q} = h^{q,p} = h^{n-p,n-q}$  we obtain a formula for the signature

$$\text{sgn}(M) = \sum_{p,q=0, 2|p+q}^{\dim(M)} (-1)^p h^{p,q}.$$

- Example: let  $n = 4$ : we sum up the terms for which  $p - q$  is even:

$$\begin{aligned} \text{sgn}(M) &= \begin{array}{ccccc} +p^{4,0} & -p^{3,1} & +p^{2,2} & -p^{1,3} & +p^{0,4} \\ & +p^{2,0} & -p^{1,1} & +p^{0,2} & \\ & & +p^{0,0} & & \end{array} \\ &= \begin{array}{ccccc} +h^{4,0} & -h^{3,1} + h^{2,0} & +h^{2,2} - h^{1,1} & -h^{1,3} + h^{0,2} & +h^{0,4} \\ & +h^{2,0} & -h^{1,1} + h^{0,0} & +h^{0,2} & \\ & & +h^{0,0} & & \end{array} \\ &= \begin{array}{ccccc} & & & +h^{4,4} & \\ & +h^{4,2} & -h^{3,3} & +h^{2,4} & \\ +h^{4,0} & -h^{3,1} & +h^{2,2} & -h^{1,3} & +h^{0,4} \\ & +h^{2,0} & -h^{1,1} & +h^{0,2} & \\ & & +h^{0,0} & & \end{array} \end{aligned}$$

- We can neglect the remaining summands with  $p + q$  odd, since  $(-1)^q h^{p,q}$  cancels with  $(-1)^p h^{q,p}$

$$\text{sgn}(M) = \sum_{p,q=0}^{\dim(M)} (-1)^p h^{p,q}$$

Further we can transform the formula:

$$\text{sgn}(M) = \sum_{p,q=0}^{\dim(M)} (-1)^q h^{p,q} = \sum_{p=0}^{\dim M} \chi(M; \Omega^p).$$

- Example: For the connected surfaces the intersection form is of the type  $(2h^{2,0} + 1, h^{1,1} - 1)$ .



## Motivation leading to the notion of Čech cohomology :

[B. V. Shabath, Introduction to complex analysis II, Chapter IV].

**9.9 Additive Cousin Problem:** find a global meromorphic function with prescribed poles.

Let  $M = \bigcup U_i$  be a covering. On each  $U_i$  there is given a meromorphic function  $f_i$ . We assume that the differences  $g_{ij} = (f_i)|_{U_i \cap U_j} - (f_j)|_{U_i \cap U_j}$  are holomorphic. Does there exist a meromorphic function  $f$  on  $M$  such that each difference  $f|_{U_i} - f_i$  is holomorphic?

**9.10 Multiplicative Cousin Problem:**

Let  $\{U_i\}_{i \in I}$  be a covering of  $M$ . On each  $U_i$  there is given a meromorphic function  $f_i$ . We assume that the quotients  $g_{ij} = \frac{(f_i)|_{U_i \cap U_j}}{(f_j)|_{U_i \cap U_j}}$  are holomorphic. Does there exist a meromorphic function  $f$  on  $M$  such that each quotient  $\frac{f|_{U_i}}{f_i}$  is holomorphic?

**9.11** The answer is in the language of Čech cohomology. For a covering  $\mathcal{U} = \{U_i\}$  the Čech complex is defined by:

$$\check{C}^k(\mathcal{U}) = \prod_{i_0 < i_1 < \dots < i_k} \mathcal{F}(U_{i_0} \cap U_{i_1} \cap \dots \cap U_{i_k}).$$

Notation: for a multiindex  $I = \{i_0 < i_1 < \dots < i_k\}$  let  $U_I = U_{i_0} \cap U_{i_1} \cap \dots \cap U_{i_k}$ . For  $\{s_I\} \in \check{C}^{k-1}(\mathcal{U})$  define the differential

$$d(\{s_I\})_J = \sum_{a=1}^k (-1)^a (s_{J \setminus j_a})|_{U_J}$$

For example

$$\begin{aligned} d(\{s_i\})_{j_0, j_1} &= (s_{j_1})|_{U_{j_0, j_1}} - (s_{j_0})|_{U_{j_0, j_1}} \\ d(\{s_{i_0, i_1}\})_{j_0, j_1, j_2} &= s_{j_1, j_2} - s_{j_0, j_2} + s_{j_0, j_1} \quad \text{restricted to } U_{j_0, j_1, j_2} \end{aligned}$$

**9.12** Čech cohomology is defined by  $\check{H}^k(\mathcal{U}; \mathcal{F}) = H^k(\check{C}^\bullet(\mathcal{U}; \mathcal{F}), d)$ .

**9.13 Additive Cousin Problem :** Let  $\mathcal{F} = \mathcal{O}_M$ , the collection of functions  $\{g_{i,j}\} \in \check{C}^1(\mathcal{U}; \mathcal{O}_M)$  satisfies the cocycle condition:

$$g_{ij} - g_{ik} + g_{jk} = 0.$$

It defines an element of Čech cohomology of the covering  $H^1(\{U_i\}; \mathcal{O}_M)$ . The cohomology class is trivial if the cocycle is a coboundary, i.e. there exists a collection of elements  $h_i \in \mathcal{O}_M(U_i)$  such that  $g_{ij} = h_j - h_i$ .

• the Cousin problem has a solution if and only if the cohomology class  $[g_{ij}] = 0$ .

Proof: If  $g_{ij} = h_j - h_i$ , then the meromorphic functions  $\tilde{f}_i = f_i + h_i$  agree at the intersections:

$$\tilde{f}_i - \tilde{f}_j = f_i + h_i - f_j + h_j \quad \text{on } U_i \cap U_j.$$

(The converse - exercise.)

**9.14** Multiplicative Cousin problem has a positive solution if the cocycle  $g_i/g_j$  defines the trivial class in  $H^1(\{U_i\}; \mathcal{O}_M^*)$ .

**9.15** Passing to a finer cover defines a map of Čech cohomology (it does not depend on inscribing function).

**9.16 Theorem:** If  $M$  is paracompact, then

$$\begin{aligned} \lim_{\mathcal{U}} \check{H}^k(\mathcal{U}; \mathcal{F}) &\simeq H^k(M; \mathcal{F}) \\ &\longrightarrow \end{aligned}$$

(The RHS is in the sense of homological algebra.)

**9.17** If the covering is acyclic (i.e.  $H^k(U_I; \mathcal{F}) = 0$  for any multiindex  $I$  and  $k > 0$ ) then

$$H^k(\{U_i\}; \mathcal{F}) \simeq H^k(M; \mathcal{F}).$$

**9.18** Sufficient conditions for being acyclic:

- For locally constant sheaves on topological spaces: if all  $U_I$  are contractible,
- For coherent sheaves in algebraic geometry: if  $U_I$  are affine,
- For coherent sheaves in analytic geometry: if  $U_I$  are Stein spaces

Definition  $U \subset M$  is Stein if:

- for any pair of points  $p, q \in U$  there exists an analytic function  $f \in \mathcal{O}_U$  such that  $f(p) \neq f(q)$ .
- (holomorphic convexity) for any compact set  $K \subset U$  the set

$$\bar{K} := \{p \in U \mid \forall f \in \mathcal{O}_U \mid f(p) \leq \sup_{q \in K} |f(q)|\}$$

is compact.

**9.19** In the Cousin problems one can pass to a finer coverings. Since  $H^1(\mathbb{P}^n; \mathcal{O}_M) = 0$ , so on  $\mathbb{P}^n$  the additive Cousin problem has always a positive solution. On curves of positive genus - not always:  $\text{genus} = \dim H^1(C; \mathcal{O}_C)$ .

## 10 Vector bundles and connection

**10.1** Let  $Vect^1(X)$  denotes the set of isomorphism classes of (topological) complex linear bundles  $X$ . Looking at the definition of Čech cohomology we discover a bijection

$$Vect^1(X) = H^1(X; C(-, \mathbb{C}^*)),$$

where  $C(-, \mathbb{C}^*)$  denotes the sheaf of continuous functions with values in  $\mathbb{C}^*$ .

- Similarly the isomorphism classes of holomorphic vector bundles over complex manifolds are identified with  $H^1(X; \mathcal{O}_X^*)$ .

**10.2** The exponential exact sequence of sheaves

$$0 \rightarrow \underline{\mathbb{Z}} \rightarrow C(-, \mathbb{C}) \xrightarrow{\exp} C(-, \mathbb{C}^*) \rightarrow 0$$

induces the map

$$c_1 : Vect^1(X) = H^1(X; C(-, \mathbb{C}^*)) \rightarrow H^2(X, \mathbb{Z}).$$

This is the first Chern class, we will give a differential definition later.

### Divisors and line bundles, [Huybrechts §2.3]

We identify holomorphic bundles with sheaves of holomorphic sections. Locally free sheaves of  $\mathcal{O}_X$ -modules are identified with holomorphic vector bundles.

**10.3** Divisor  $D = \sum a_i D_i$  is a formal combination of codimension 1 indecomposable subvarieties (we assume that  $X$  is an analytic manifold).

- We define a restriction of divisors to open sets:  $D|_U = \sum_{D_i \cap U \neq \emptyset} a_i (D_i \cap U)$
- $D$  is an effective divisor iff all  $a_i \geq 0$ , we write  $D \geq 0$ .

**10.4** A meromorphic function defines a principal divisor  $\text{div}(f) = \text{zeros}(f) - \text{poles}(f)$ .

**10.5** Any divisor  $D$  defines a line bundle  $\mathcal{O}_X(D)$ , viewed as a subsheaf of the sheaf  $Mero_X$  of meromorphic functions: for each open  $U \subset X$

$$\mathcal{O}_X(D)(U) = \{f \in Mero_X(U) : \text{div}(f) + D|_U \text{ is effective in } U\}$$

- If  $D_1 = D_2 + \text{div}(g)$ , where  $g$  is a global meromorphic function, then  $\mathcal{O}_X(D_1) \simeq \mathcal{O}_X(D_2)$ . The multiplication by  $g$  defines an isomorphism.
- We have an injection

$$\{\text{Divisors}\} / \{\text{Principal divisors}\} \hookrightarrow \{\text{Holomorphic Line Bundles}\}.$$

The image consists of line bundles admitting a meromorphic section.

- Suppose  $s : X \dashrightarrow L$  is a meromorphic section. Define  $D = \text{zeros}(s) - \text{poles}(s)$ . Then  $L \simeq \mathcal{O}_X(D)$
- If  $L \rightarrow X$  is an algebraic bundle, then it admits a meromorphic section.

**10.6** Example: the tautological bundle over  $\mathbb{P}^1$ .

On  $U_0 = \{z_0 \neq 0\}$  we have a section  $s_0([1 : z]) = (1, z)$ , on  $U_1 = \{z_1 \neq 0\}$  we have a section  $s_1([w : 1]) = (w, 1)$ . These sections do not vanish, so they define local trivializations. The transition function  $g_{1,0}s_0 = s_1$  satisfies

$$g_{1,0}(z)(1, z) = (1/z, 1).$$

Hence

$$g_{1,0} : U_0 \cap U_1 = \mathbb{C}^* \rightarrow GL_1(\mathbb{C}) = \mathbb{C}^*,$$

$$g_{0,1}(z) = z^{-1}.$$

- The section  $s_0$  has pole at  $\infty$ , hence the tautological bundle is isomorphic to  $\mathcal{O}_{\mathbb{P}^1}(D)$ , where  $D = -\{[0 : 1]\}$ . Equally well we could have  $D = -\{[1 : 0]\}$  or any other point.

**10.7** Taking the transition function  $g_{1,0}(z) = z^k$  we obtain  $\mathcal{O}_{\mathbb{P}^1}(k)$

**10.8** Example:  $\mathcal{O}_{\mathbb{P}^n}(kH)$ , where  $H \simeq \mathbb{P}^{n-1}$  is the divisor at infinity. If  $k \geq 0$  the global sections  $H^0(\mathbb{P}^n; \mathcal{O}_{\mathbb{P}^n}(kH))$  are naturally identified with  $\mathbb{C}[z_1, z_2, \dots, z_n]_{\deg \leq k} \simeq \mathbb{C}[z_0, z_1, \dots, z_n]_{\deg = k}$  and

$$\mathcal{O}_{\mathbb{P}^n}(kH) \simeq (\text{tautological}^*)^{\otimes k} =: \text{tautological}^{\otimes -k}.$$

The only section for  $k < 0$  is 0 and

$$\mathcal{O}_{\mathbb{P}^n}(kH) \simeq \text{tautological}^{\otimes -k}.$$

**10.9** The bundle  $\mathcal{O}_{\mathbb{P}^n}(kH) \simeq \text{tautological}^{\otimes -k}$  is denoted  $\mathcal{O}_{\mathbb{P}^n}(k)$ .

- If  $Y$  is a hypersurface in  $\mathbb{P}^n$  of degree  $d$ , then  $\mathcal{O}_{\mathbb{P}^n}(Y) \simeq \mathcal{O}_{\mathbb{P}^n}(d)$ .

## Connection for a vector bundle over $C^\infty$ -manifold

**10.10** Connection is a linear map  $\nabla : C^\infty(X; E) \rightarrow C^\infty(X; T_X^* \otimes E) =: A_X^1(E)$  satisfying the Leibniz rule

$$\nabla(fs) = df \otimes (s) + fs.$$

**10.11** Let  $\nabla$  and  $\nabla'$  be two connections. The difference  $\nabla - \nabla'$  is  $A_X^1$  linear. [Huybrechts 4.2.3]

- Locally, every connection is of the form  $\nabla = d + a$ , where  $a \in A^1(X, \text{End}(E))$ .
- If  $\nabla$  is a connection and  $a \in A^1(X, \text{End}(E)) = C^\infty(X, T^*X \otimes \text{End}(E))$ , then  $\nabla + a$  is a connection.
- Affine combination of connections  $t\nabla_1 + (1-t)\nabla_2$  is a connection.
- Applying a partition of unity associated to the trivializing atlas of  $E$  we glue together local connections and obtain a global one.
- The space of connections is isomorphic to  $A^1(X, \text{End}(E))$ . (But no connection is distinguished.)

## Connections concordant with structures

**10.12** Suppose  $E$  is a hermitian bundle. A connection is Hermitian if

$$d \langle s_1, s_2 \rangle = \langle \nabla s_1, s_2 \rangle + \langle s_1, \nabla s_2 \rangle.$$

(again the Leibniz formula) [Huybrechts 4.2.9]

**10.13** Let  $V$  be a Hermitian vector space. By  $\text{End}(V, h)$  denote the endomorphism  $a$  satisfying

$$\langle a(v), w \rangle + \langle v, a(w) \rangle = 0.$$

If  $V = \mathbb{C}^n$  with the standard hermitian product, then  $\text{End}(V, h) = \mathfrak{u}_n = \{A \in M_{n \times n}(\mathbb{C}) \mid A + \bar{A}^T = 0\}$ .

- For a Hermitian vector bundle  $\text{End}(V, h)$  is a real vector bundle of the dimension  $= \dim(\mathfrak{u}_{\text{rk } E})$ .
- As before we prove that the space of Hermitian connection is a real vector space isomorphic to  $A^1(X, \text{End}(E, h))$ . (But no connection is distinguished.)

**10.14** If  $\text{rk } E = 1$ . Then  $\text{End}(E, h) \simeq \mathbb{R}$ .

**10.15** Suppose  $X$  is a complex manifold,  $E$  is a holomorphic bundle (the transition functions are holomorphic). Let  $A^k(X, E) = \Gamma(A_X^k \otimes E)$ ,  $A^k(X, E) = \bigoplus_{p+q=k} A^{p,q}(X, E)$ . The operator  $\bar{\partial}$  is well defined

$$\bar{\partial}_E : A^{p,q}(X, E) \rightarrow A^{p,q+1}(X, E).$$

Warning: the operator  $\partial$  does not commute with the transition functions. Thus  $\partial_E$  is not defined, unless the transition functions are locally constant.

**10.16** The connection decomposes into components  $\nabla^{1,0} + \nabla^{0,1}$ . We say that  $\nabla$  is compatible with the complex structure if  $\nabla^{0,1} = \bar{\partial}_E$ .

**10.17** The space of connections compatible with complex structure is isomorphic to  $A^{1,0}(X, \text{End}(E))$ .

**10.18** Theorem [Huybrechts 4.2.14]: For a Hermitian holomorphic bundle there exists exactly one connection compatible with the complex structure.

- In local coordinates: let  $H$  be the matrix of the Hermitian product,  $\nabla = d + A$ , (we identify  $A$  locally with a matrix, we call it connection matrix)

$$A \in M_{n \times n}(A^{1,0}(X)), \quad H \in M_{n \times n}(C^\infty(X)), \quad \bar{H} = H^T, \quad n = \text{rk}(E).$$

- the Hermitian condition reads

$$dH = A^T H + H \bar{A}.$$

Hence

$$\partial H = A^T H,$$

so

$$A = \bar{H}^{-1} \partial(\bar{H}).$$

- If  $n = 1$ ,  $H = [h]$ . then  $a = \partial \log(h)$ .

**10.19** Example  $L = \mathcal{O}(-1)$  on  $\mathbb{P}^n$ , i.e. the tautological bundle,  $L \subset \mathbb{C}^{n+1} \times \mathbb{P}^n$  has the induced Hermitian structure from the trivial bundle  $\mathbb{C}^{n+1}$ . The connection form

$$A = \partial \log(\|s\|^2),$$

where  $s$  is any section (trivialization) of  $L$ .

- For example on the chart  $\{z_0 \neq 0\} \simeq \mathbb{C}^n$  there is a section

$$s([1 : z_1 : \cdots : z_n]) = (1 : z_1 : \cdots : z_n),$$

the differential

$$F_\nabla = d\partial \log(1 + \|z\|^2) = -\partial \bar{\partial} \log(1 + \|z\|^2),$$

is called the curvature.

- Note:

$$\frac{i}{2\pi} F_\nabla = -\omega_{FS}.$$

## 11 Chern classes

**11.1** We extend the connection using Leibniz formula to obtain the operator  $\nabla_E : A^k(X, E) \rightarrow A^{k+1}(X, E)$ .

**11.2** Theorem: The curvature  $F_\nabla = \nabla^2 : A^0(E) \rightarrow A^2(E)$  is  $A^0(X)$ -linear, hence it defines a section of the bundle  $\Lambda^2 T^*X \otimes \text{End}(E)$ .

**11.3** Locally in the matrix notation

$$F_\nabla = dA + A \wedge A \in M_{n \times n}(A^2(X)).$$

**11.4** For a line bundle  $E = L = \mathbb{C} \times X$ : we have  $\text{End}(L) = \mathbb{C}$  and  $A \wedge A = 0$  (since  $A$  is a  $1 \times 1$  matrix). Then  $H = [h]$ ,  $h : X \rightarrow \mathbb{R}$

$$F_\nabla = dA = d\partial \log(h) = \bar{\partial} \partial \log(h).$$

**11.5** Example  $L = \mathcal{O}(-1)$  on  $\mathbb{P}^n$ , i.e. the tautological bundle,  $L \subset \mathbb{C}^{n+1} \times \mathbb{P}^n$  has the induced Hermitian structure from the trivial bundle  $\mathbb{C}^{n+1}$ :

$$F_\nabla = d\partial \log(\|s\|^2) = -\partial \bar{\partial} \log(\|s\|^2),$$

where  $v$  is any section of  $L$ ,

$$\frac{i}{2\pi} F_\nabla = -\omega_{FS}.$$

- For example on the chart  $\{z_0 \neq 0\} \simeq \mathbb{C}^n$  there is a section

$$s([1 : z_1 : \cdots : z_n]) = (1 : z_1 : \cdots : z_n),$$

hence

$$F_\nabla = -\partial \bar{\partial} \log(1 + \|z\|^2).$$

- Note:  $c_1(\mathcal{O}(-1)) = -[\omega_{FS}]$ .

**11.6** Connection on  $E$  induces a connection on  $\text{End}(E)$

$$(\nabla f)(s) := \nabla(f(s)) - f\nabla s = [\nabla, f]s.$$

In particular we can apply  $\nabla$  to  $F_\nabla$ .

**11.7** Bianchi identity:

$$\boxed{\nabla(F_\nabla) = 0 \in A^3(X, \text{End}(E))},$$

because  $[\nabla, \nabla \circ \nabla] = 0$ .

- Locally for  $\nabla = d + A$  we have

$$dF_\nabla = d(dA + A \wedge A) = dA \wedge A - A \wedge dA = [dA, A] = [F_\nabla, A].$$

Hence

$$0 = \nabla(F_\nabla) = dF_\nabla + [A, F_\nabla].$$

We obtain a formula for the differential

$$\boxed{dF_\nabla = [F_\nabla, A]}.$$

## Differential definition of Chern classes

Huybrechts §4.4

**11.8 Theorem:** For any polynomial map  $P : \text{End}(\mathbb{C}^n) \rightarrow \mathbb{C}$  which is invariant with respect to conjugation the form  $P(\nabla_E^2) \in A^{2\deg(P)}(X)$  is closed.

• Lemma (see Milnor-Stasheff, Appendix C, p.297) For  $X = (x_{ij})_{i,j}$  define the matrix  $P'(X) = (\frac{\partial P}{\partial x_{ji}})_{i,j}$  (note, that the indices  $i, j$  are exchanged). We have:

(1)  $dP(X) = \text{tr}(P'(X) \cdot dX)$ .

(2) if  $P$  is Ad-invariant, then the matrices  $P'(X)$  and  $X$  commute.

Proof:

ad (1) easy

ad (2)  $P((I + tE_{ij})X) = P(X(I + tE_{ij}))$ , hence

$$\sum_k x_{i,k} \frac{\partial P}{\partial x_{jk}} = \sum_k \frac{\partial P}{\partial x_{ki}} x_{k,j}$$

• Proof of theorem:

$$\begin{aligned} dP(F_\nabla) &\stackrel{(1)}{=} \text{tr}(P'(F_\nabla)dF_\nabla) = \text{tr}(P'(F_\nabla)[F_\nabla, A]) = \text{tr}(P'(F_\nabla) \wedge F_\nabla \wedge A - P'(F_\nabla) \wedge A \wedge F_\nabla) = \\ &\stackrel{(2)}{=} \text{tr}(F_\nabla \wedge (P'(F_\nabla) \wedge A) - (P'(F_\nabla) \wedge A) \wedge F_\nabla) = \text{tr}([F_\nabla, P'(F_\nabla) \wedge A]) = 0. \end{aligned}$$

**11.9 Remark:** the map

$$\mathbb{C}[M_{n \times n}(\mathbb{C})]^{GL_n} \rightarrow \mathbb{C}[\text{diagonal matrices}]^{\Sigma_n} = \mathbb{C}[\sigma_1, \sigma_2, \dots, \sigma_n]$$

is an isomorphism. If  $P$  is Ad-invariant, then it can be expressed by the coefficients of the characteristic polynomial. Equivalently,  $P(A)$  is a symmetric function in eigenvalues of  $A$ .

**11.10** The 2-form  $P(F_\nabla)$  defines a cohomology class, which does not depend on the connection (dowód TBA).

• For  $P = (\frac{i}{2\pi})^k \sigma_k$ , ( $(-1)^k \sigma_k$  is  $(\text{rk} E - k)$ -th coefficient of the characteristic polynomial) the resulting forms represent the Chern classes.

## The first Chern class $c_1(L)$ of a line bundle - various constructions

• Axiomatic definition, see Milnor-Stasheff:

<p>a) <math>c_1 : \text{Vect}^1(-) \rightarrow H^2(-, \mathbb{Z})</math> is a natural transformation of functors <math>\text{Top} \rightarrow \text{GrAb}</math></p> <p>b) <math>c_1(L_1 \otimes L_2) = c_1(L_1) + c_1(L_2)</math></p> <p>c) <math>c_1(\mathcal{O}_{\mathbb{P}^1}(1)) =</math> the distinguished generator <math>[pt] \in H^2(\mathbb{P}^1)</math></p>
--

• The identification  $\text{Vect}^1(X) = [X, \mathbb{P}^\infty] = [X, K(\mathbb{Z}, 2)] = H^2(X; \mathbb{Z})$ ,

where  $[-, -]$  denotes the set of homotopy classes of maps.

• Via the differential in the long exact sequence of cohomologies associated to the short exact sequence of sheaves

$$0 \rightarrow 2\pi i\mathbb{Z} \rightarrow C(X, \mathbb{C}) \xrightarrow{\exp} C(X, \mathbb{C}^*) \rightarrow 0$$

$$0 = H^1(X, C(X, \mathbb{C})) \rightarrow H^1(X, C(X, \mathbb{C}^*)) \xrightarrow{\sim} H^2(X; 2\pi i\mathbb{Z}) \rightarrow H^2(X, C(X, \mathbb{C})) = 0$$

Note, that we have an identification  $\text{Vect}^1(X) = \check{H}^1(X, H^1(X, C(X, \mathbb{C}^*)))$ .

• via the obstruction theory: the obstruction to the existence of a nonzero section belongs to

$$H^2(X; \pi_1(\mathbb{C}^*)) \simeq H^2(X; \mathbb{Z})$$

•  $c_1(L) = [\text{zero section}] \in H^2(L) \simeq H^2(X)$

• a definition via connection (when  $X$  is a manifold) t.j.w.  $\frac{i}{2\pi}[F_\nabla] = \frac{i}{2\pi}[\partial\bar{\partial}\log h] \in H^2(X; \mathbb{C})$ , [Huybrechts §4].

## Generalities about characteristic classes for higher rank bundles

- Let

$$Vect^n(X) = \{\text{isomorphism classes of } n\text{-dimensional complex vector bundles over } X\}$$

- Def: a characteristic class on  $n$ -dimensional bundle is a transformation of functors  $hTop \rightarrow Sets$

$$Vect^n(-) \longrightarrow H^*(-).$$

Since  $Vect^n(-)$  is representable,

$$Vect^n(X) = \{\text{homotopy classes } f : X \rightarrow Grass_n(\mathbb{C}^\infty)\}$$

for finite  $CW$ -complexes, by Yoneda lemma

$$\{\text{Characteristic classes of } n\text{-bundles}\} = H^*(Grass_n(\mathbb{C}^\infty)) = \mathbb{Z}[c_1, c_2, \dots, c_n].$$

- More generally, for a compact Lie group (or a reductive algebraic group)  $G$ , for cohomology with coefficients in  $\mathbb{C}$ :

Let  $Bun^G(X)$  be the set of isomorphism classes of  $G$ -bundles over  $X$ . This functor  $hTop \rightarrow Set$  is representable by  $BG$ , thus

$$\{\text{Characteristic classes of } G\text{-bundles}\}_{\mathbb{C}} = H^*(BG; \mathbb{C}) = \mathbb{C}[\mathfrak{g}]^G = \mathbb{C}[\mathfrak{t}]^W.$$

by Borel theorem. Here  $\mathfrak{g}$  is the Lie algebra of  $G$ ,  $\mathfrak{t}$  the Lie algebra of the maximal torus, and  $W = NT/T$  is the Weyl group.

- For  $G = GL_n$  we have

$$\mathbb{C}[\mathfrak{t}]^W = \mathbb{C}[t_1, t_2, \dots, t_n]^{\Sigma_n},$$

the ring of symmetric functions in  $n$  variables.

### 11.11 Axioms of Chern classes

$$c(E) = 1 + c_1(E) + c_2(E) + \dots + c_r(E), \quad r = rk(E), \quad c_k(E) \in H^{2k}(X; \mathbb{Z}).$$

- 1) Functoriality:  $c(-)$  is a transformation of functors

$$Vect^r(-) \rightarrow H^*(-; \mathbb{Z}),$$

- 2) Whitney formula:

$$c(E_1 \oplus E_2) = c(E_1) \cup c(E_2),$$

where  $\cup$  denotes product in cohomology, sometimes written simply as  $\cdot$ .

- 3) Normalization:

$$c(\mathcal{O}_{\mathbb{P}^1}(1)) = 1 + [pt],$$

where  $[pt]$  is the generator of  $H^2(\mathbb{P}^1; \mathbb{Z})$ , which agrees with the complex orientation. (In de Rham cohomology  $[pt] = [\omega_{FS}]$ ).

- By splitting principle we can assume that a vector bundle is a sums of line bundles. The cost is that we replace the base  $X$  by  $Fl(E)$ , the bundle of flags over  $X$ , which is harmless, since it is mono on cohomology. Topologically every line bundle can be pulled back from  $\mathbb{P}^\infty$  (which has the same cohomology  $H^2$  as  $\mathbb{P}^1$ ). Hence the axioms determine  $c(E)$ .

## 12 Chern classes and others

The total  $c(E)$  is associated to the Ad-invariant (nonhomogeneous) polynomial  $P : X \mapsto \det(X + I)$

**12.1** Let  $f : Y \rightarrow X$  be a  $C^\infty$ -map,  $E \rightarrow X$  a complex bundle with a connection  $\nabla$ . The **pull back**: if locally  $\nabla = d_X + A$ , then  $f^*\nabla = d_Y + f^*A$  - (Huybrechts 4.2.6.v)

$$\begin{array}{ccccc} \text{End}(f^*E) \otimes T^*Y|_{f^{-1}U} & \longleftarrow & \text{End}(f^*E) \otimes f^*T^*X|_{f^{-1}U} & \longrightarrow & \text{End}(E) \otimes T^*X|_U \\ & \nwarrow f^*A & \downarrow & & \nearrow A \uparrow \downarrow \\ & & f^{-1}(U) & \longrightarrow & U \subset X \end{array}$$

**12.2** Let  $P$  be an Ad-invariant polynomial. The class of  $P(F_\nabla)$  in  $H^*(X)$  does not depend on the connection. (Assumption:  $X$  is a real manifold.)

Proof: for two connections  $\nabla_0, \nabla_1$  define a connection on  $X \times \mathbb{R}$  by the formula  $\tilde{\nabla} = t p^* \nabla_1 + (1-t) p^* \nabla_0$ , where  $p : X \times \mathbb{R} \rightarrow X$  is the projection. Inclusions  $i_0, i_1 : X \rightarrow X \times \mathbb{R}$  on submanifolds  $t = 0$  and  $t = 1$  are homotopic, so  $[P(F_{\nabla_1})] = i_1^* P([F_{\tilde{\nabla}}]) = i_0^* [P(F_{\tilde{\nabla}})] = [P(F_{\nabla_0})]$ .

**12.3** Verification of axioms:

- 1) Functoriality (the connection can be pulled back)
- 2) Whitney formula

$$c(E_1 \oplus E_2) = c(E_1)c(E_2),$$

Let connection on  $E_1 \oplus E_2$  be of the product form  $\nabla(s_1, s_2) = (\nabla_1 s_1, \nabla_2 s_2)$ . Then

$$F_\nabla = F_{\nabla_1} \oplus F_{\nabla_2} \in (\text{End}(E_1) \oplus \text{End}(E_2)) \otimes A^2(X) \subset \text{End}(E_1 \oplus E_2) \otimes A^2(X).$$

- 3) Normalization  $c_1(\mathcal{O}(-1)) = -[\omega_{FS}]$ , by the definition of the Fubini-Study form.

**12.4** Remark: The differential forms obtained by the above constructions are integral (i.e. come from  $H^*(X; \mathbb{Z})$ ).

**12.5** Suppose  $X$  is complex manifold,  $L$  a holomorphic line bundle with a Hermitian structure. Then  $F_\nabla = dA = \bar{\partial}\partial \log h$  is a (1,1)-form.

- If  $X$  is Kähler manifold, then  $c_1(L) \in H^{1,1}(X) \cap \text{image}(H^2(X; \mathbb{Z}))$ .

**12.6** Theorem: Let  $X$  be a Kähler manifold,  $E$  a holomorphic bundle, then  $c_k(E) \in H^{2k}(X; \mathbb{C})$  is represented by a  $(k, k)$ -form  $(\frac{i}{2\pi})^k \sigma_k(F_\nabla)$ .

- Splitting principle: Let  $p : Fl(E) \rightarrow X$  be a bundle of flag spaces over  $X$ .

$$Fl(E) = \{(x, V_1, V_2, \dots, V_{\text{rk}E}) \mid x \in X, V_i \subset E_x, \dim(V_i) = i, V_i \subset V_{i+1}\}.$$

Let  $L_i = V_i/V_{i-1}$ . The Hermitian structure defines an isomorphism  $V_i = L_i \oplus V_{i-1}$ . (Note: This isomorphism is not holomorphic.)

Topologically  $p^*E = \bigoplus_{i=1}^{\text{rk}E} L_i$ . The Chern classes are topological invariants, hence

$$c_\bullet(p^*E) = \prod_{i=1}^{\text{rk}E} (1 + c_1(L_i)).$$

Each  $c_1(L_i)$  is of the type (1,1) thus  $c_k(p^*E)$  is of the type  $(k, k)$ .

Fact:  $p^* : H^*(X) \hookrightarrow H^*(Fl(E))$  is a monomorphism. Moreover it preserves types. Conclusion:  $c_k(E)$  is of the type  $(k, k)$ .

**12.7** Huybrechts 4.2.18: in general one can define Atiyah class  $A(E) \in H^1(X; \Omega_X^1 \otimes \text{End}(E))$ , which agree with  $\frac{1}{2\pi i} F_\nabla \in A^2(X; \text{End}(E))$ .



## Other classes

**12.8** Chern character. Let  $P \in \mathbb{C}[[M_{n \times n}]]$  be given by the formula:

$$P(B) = \sum_{k=0}^{\infty} \frac{\text{tr}(B^k)}{k!},$$

where  $B = \frac{i}{2\pi} F_{\nabla}$ . In terms of symmetric functions

$$P(t_1, t_2, \dots, t_n) = \sum_{k=0}^{\infty} \sum_{i=1}^n \frac{t_i^k}{k!} = \sum_{k=0}^{\infty} e^{t_i}.$$

The resulting characteristic class is denoted by  $ch(E)$ .

- Properties:

$$ch(E_1 \oplus E_2) = ch(E_1) + ch(E_2)$$

$$ch(E_1 \otimes E_2) = ch(E_1) \cup ch(E_2)$$

The second identity follows from  $e^{a+b} = e^a e^b$ .

**12.9** In general having a formal power series  $f[[x]]$  we define an additive characteristic class satisfying:

- $a_f(L) = f(c_1(L))$  for a line bundle  $L$
- $a_f(E_1 \oplus E_2) = a_f(E_1) + a_f(E_2)$

**12.10** Example: if  $f[[x]] = e^x$ , then  $a_f(E) = ch(E)$ .

- To express the homogeneous components of  $ch(E)$  assume that  $E$  is a sum of line bundles  $L_i$ , let  $x_i = c_1(L_i)$
- $ch(E)_{(0)} = \text{rk} E$
- $ch(E)_{(1)} = x_1 + x_2 + \dots + x_n = c_1(E)$
- $ch(E)_{(2)} = \frac{1}{2}(x_1^2 + x_2^2 + \dots + x_n^2) = \frac{1}{2}(x_1 + x_2 + \dots + x_n)^2 - \sum x_i x_j = \frac{1}{2}c_1^2(E) - c_2(E)$

**12.11** For a formal power series  $f[[x]]$  we define a multiplicative characteristic class satisfying:

- $m_f(L) = f(c_1(L))$  for a line bundle  $L$
- $m_f(E_1 \oplus E_2) = m_f(E_1) \cup m_f(E_2)$
- Example: if  $f[[x]] = 1 + x$ , then  $m_f(E) = c_{\bullet}(E)$ .

**12.12** Todd class: Let

$$f[[x]] = \frac{x}{1 - e^{-x}} = \frac{x}{x - x^2/2 + x^3/6 \dots} = 1 + \frac{x}{2} + \frac{x^2}{12} - \frac{x^4}{720} + \frac{x^6}{30240} + \dots$$

(the coefficients are  $\pm \frac{\text{Bernoulli number}}{n!}$ )

- $td(E)_{(0)} = 1$
- $td(E)_{(1)} = \frac{1}{2}c_1(E)$
- $td(E)_{(2)} = \frac{x_1^2}{12} + \frac{x_1 x_2}{4} + \frac{x_2^2}{12} = \frac{1}{12}(c_1^2(E) + c_2(E))$  (for degrees  $\leq 2$  it is enough to perform computation for  $\text{rk} E = 2$ )
- Alternative description:  $td(E)$  is associated to the function on matrices  $B \mapsto \frac{\det(B)}{\det(id - e^{-B})}$ .

**12.13** Hirzebruch-Riemann-Roch (Huybrechts 5.1.1) Let  $E$  be a holomorphic vector bundle on a compact manifold  $X$ . Then

$$\chi(X; E) = \int_X td(TX) \cup ch(E).$$

**12.14** Essentially it is enough to check the equality for  $X = \mathbb{P}^n$ ,  $E = \mathcal{O}(k)$ .

$$LHS = \dim(\mathbb{C}[x_0, x_1, \dots, x_n]_k) = \binom{n+k}{k}$$

- Lemma: Euler exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n} \rightarrow \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus n+1} \rightarrow T\mathbb{P}^n \rightarrow 0.$$

Hence

$$td(T\mathbb{P}^n) = \left( \frac{h}{1-e^{-h}} \right)^{n+1},$$

where  $h = [\omega_{FS}] \in H^2(\mathbb{P}^n)$ .

•

$$\begin{aligned} RHS &= \int_{\mathbb{P}^n} \left( \frac{h}{1-e^{-h}} e^{kh} \right)^{n+1} = \\ &= \left[ \left( \frac{h}{1-e^{-h}} \right)^{n+1} e^{kh} \right]_{\text{coef of } h^n} = Res_{h=0} \left( \frac{e^{kh}}{(1-e^{-h})^{n+1}} \right) = Res_{h=0} \frac{e^{(k+n+1)h}}{(e^h - 1)^{n+1}} = \dots \end{aligned}$$

Let  $u = e^h - 1$ ,  $du = e^h dh$

$$\dots = Res_{u=0} \frac{(u+1)^{n+k}}{u^{n+1}} = [(u+1)^{n+k}]_{\text{coef of } u^n}$$

**12.15** Exercise:  $X$  = hypersurface of degree  $d$  in  $\mathbb{P}^n$ ,  $E = \mathcal{O}(k)$ :

$$td(TX) = td(T\mathbb{P}^n)/td(\nu_X) = \left( \frac{h}{1-e^{-h}} \right)^{n+1} \frac{1-e^{-dh}}{dh} = \frac{h^n}{d} \frac{1-e^{-dh}}{(1-e^{-h})^{n+1}}$$

...

**12.16** If  $\dim X = 1$ , suppose  $L$  is of degree  $d$ , i.e.  $c_1(L) = d[pt]$  then

$$\chi(X; L) = [(1 + c_1(TX)/2)(1 + c_1(L))]_{(1)} = \deg(c_1(TX)/2 + c_1(L)) = \frac{1}{2}\chi_{top}(X) + d = 1 - \text{genus} + d.$$

**12.17** If  $\dim X = 2$ ,  $L = \mathcal{O}(D)$ , then  $c_1(L) = [D]$ . Let  $c_i = c_i(TX)$

$$\begin{aligned} \chi(X; L) &= [(1 + c_1/2 + \frac{1}{12}(c_1^2 + c_2)(1 + D + D^2/2))]_{(2)} = \deg(\frac{1}{12}(c_1^2 + c_2) + \frac{c_1 \cup D + D^2}{2}) \\ &= \chi(X; \mathcal{O}_X) + \frac{c_1 \cdot D + D^2}{2} \end{aligned}$$

Using a common notation in algebraic geometry  $c_1 = -K_X$

$$\chi(X; L) = \chi(X; \mathcal{O}_X) + \frac{(D - K_X) \cdot D}{2}.$$

or with  $p_a = -\dim H^1(X; \mathcal{O}_X) = \chi(X; \mathcal{O}_X) - 1$  (arithmetic genus)

$$\chi(X; L) = 1 + p_a + \frac{(D - K_X) \cdot D}{2}.$$

**12.18** Hirzebruch class: Let

$$f_y(x) = x \frac{1 + ye^{-x}}{1 - e^{-x}} = (1+y) \frac{x}{1 - e^{-x}} - xy = (1+y) + \frac{1}{2}(1-y)x + \frac{1+y}{12}x^2 - \frac{1+y}{720}x^4 + \frac{1+y}{30240}x^6 + \dots$$

Here  $y$  is a parameter or a free variable.

- Exercise by Hirzebruch-Riemann-Roch (Huybrechts Cor. 5.1.4)

$$\int_X \text{Hirzebruch class} = \sum_{p=0}^n \chi(X, \Omega_X^p) y^p = \sum_{p,q} h^{p,q} y^p.$$

- For  $y = -1$  we obtain  $\chi_{top}(X)$  the topological Euler characteristic,  
For  $y = 0$  we obtain  $Td(X) = \chi(X, \mathcal{O}_X)$  Todd genus  
For  $y = 1$  we obtain the signature.

## 13 Positive line bundles

### 13.1 Riemann-Roch

$$\chi(X; E) = \int_X td(TX) \cup ch(E).$$

Holomorphic invariant = Topological data.

- Goal: compute  $\dim \Gamma(X, E) = \dim H^0(X; E)$ .
- In general it is not possible (by topological data).
- If  $H^k(X; E) = 0$  for  $k > 0$ , then  $\dim \Gamma(X, E) = \chi(X; E)$ .
- Hence importance of *vanishing theorems*.

**We concentrate on linear bundles**

### 13.2 Linear algebra:

$$\{1_{\frac{1}{2}} - \text{linear forms on } V\} \leftrightarrow \Lambda^2 V_{\mathbb{R}}^* \cap \Lambda^{1,1} V^* \subset \Lambda^2(V^* \otimes_{\mathbb{R}} \mathbb{C}).$$

**13.3** We say that  $\omega \in A^{1,1}(X) \cap A^2(X; \mathbb{R}) \subset A^2(X; \mathbb{C})$  is positive, if there exists a hermitian product such that  $\omega$  is equal to minus its imaginary part  $\langle\langle x, y \rangle\rangle = \langle x, y \rangle - i\omega(x, y)$ . If locally in some coordinates the hermitian product is given by a matrix  $H = [h_{i,j}]$ , then

$$\omega = \frac{i}{2} \sum h_{ij} dz_i \wedge \bar{d}z_j.$$

**13.4** A linear bundle is positive if it admits a connection  $\nabla$  such that  $\frac{i}{2\pi} F_{\nabla}$  is a positive (1,1)-form.

**13.5** Example of positive bundles:  $\mathcal{O}_{\mathbb{P}^n}(k)$ ,  $k > 0$ .

**13.6** Denote  $\Omega_X^{\dim X}$  by  $K_X$ , call it the canonical bundle/divisor.

### 13.7 Kodaira(-Nakano) Vanishing theorem

[Huybrechts Proposition 5.2.2, Griffiths-Harris p 154.]

If  $L$  is positive, then  $H^i(X; K_X \otimes L) = 0$  for  $i > 0$ .

(In algebraic geometry notation  $L = \mathcal{O}(D)$ . The vanishing theorem reads  $H^i(X; K_X + D) = 0$ .)

[Dowód na końcu w §14]

**13.8** Corollary: Assume  $L$  is positive. For any line bundle  $L'$  we have vanishing  $H^i(X; L^{\nu} \otimes L') = 0$  for  $i > 0$ ,  $\nu \gg 0$ .

•

$$L^{\nu} \otimes L' = K_X \otimes (K_X^{-1} \otimes L^{\nu} \otimes L').$$

The bundle  $K_X^{-1} \otimes L^{\nu} \otimes L'$  has the connection form equal to

$$-F_{\nabla_{K_X}} + \nu F_{\nabla_L} + F_{\nabla_{L'}}.$$

For sufficiently large  $\nu$  it is positive.

- If  $K_X^{-1}$  is positive, then vanishing holds already for  $\nu = 1$ . (Fano manifold, e.g.  $\mathbb{P}^n$ , Grassmannians, flag varieties.)

**13.9** If  $L$  is generated by global holomorphic sections (we say „globally generated”) iff for each  $x \in X$  there exists a section  $s \in H^0(X; L)$ , such that  $s(x) \neq 0$ .

**13.10** Let  $s_0, s_2, \dots, s_r$  be the basis of  $H^0(X; L)$ . Define  $\phi : X \rightarrow \mathbb{P}^r$  by

$$x \mapsto [s_0 : s_1 : \dots : s_r].$$

- Coordinate-free construction: If  $L$  is globally generated, then for  $x \in X$  define a function on the space of the global sections  $H^0(X; L)$

$$\Phi(x) \in \text{Hom}(H^0(X; L), L_x) \xrightarrow{\text{up to a scalar}} H^0(X; L)^*,$$

$$\Phi(x) : s \mapsto s(x) \in L_x \simeq \mathbb{C}.$$

- We obtain a natural map

$$\phi : X \rightarrow \mathbb{P}(H^0(X; L)^*).$$

- Then  $L = \phi^*(\mathcal{O}(1))$  (by tautological identification  $H^0(\mathbb{P}(V^*); \mathcal{O}(1)) \simeq V$ ). Hence  $L$  is „nonnegative”, i.e.  $L$  admits a connection such that the associated Hermitian form is nonnegative semi-definite.
- This property is preserved by pull-backs.

### 13.11 Kodaira embedding theorem.

[Huybrechts Proposition §5.3, Griffiths-Harris p 176.]

If a bundle  $L$  is positive, then for  $\nu \gg 0$  the tensor power  $L^\nu$  is generated by global sections and the natural map  $X \rightarrow \mathbb{P}(H^0(X; L^\nu)^*)$  is an embedding. (We only assume that,  $X$  is a compact analytic complex manifold, and as a corollary from GAGA we obtain that  $X$  is algebraic.)

- Steps of the proof:

a) assume that  $\phi_{L^\nu}$  is well defined, i.e. the base locus of  $L^\nu$  is empty:

$$\forall x \in X \quad H^0(X; L^\nu) \rightarrow L_x^\nu = H^0(X; L^b \otimes \mathcal{O}_X/m_x)$$

b)  $\phi_{L^\nu}$  separates the points: suppose for  $x \in X$  the restriction  $H^0(X; L^\nu) \rightarrow L_x^\nu \oplus L_y^\nu$  is surjective.

(note b)  $\Rightarrow$  a) )

b') Equivalently: the restriction

$$H^0(\tilde{X}; \tilde{L}^\nu) \rightarrow H^0(E; \tilde{L}_{|E}^\nu) = H^0(\tilde{X}; \tilde{L}^\nu \otimes \mathcal{O}_{\tilde{X}}/I_E)$$

is surjective, where  $\tilde{X} = Bl_x Bl_y X$ ,  $\tilde{L}$  is the pull-back of  $L$  to  $\tilde{X}$ ,  $E$  the sum of the exceptional divisors,  $I_E \simeq \mathcal{O}(-E)$  the ideal sheaf of  $E$ . The restriction map is a part of an exact sequence

$$\rightarrow H^0(\tilde{X}; \tilde{L}^\nu) \rightarrow H^0(\tilde{X}; \tilde{L}^\nu \otimes \mathcal{O}_{\tilde{X}}/I_E) \rightarrow H^1(\tilde{X}; \tilde{L}^\nu \otimes \mathcal{O}(-E)) \rightarrow$$

obtained from

$$0 \rightarrow \mathcal{O}_{\tilde{X}}(-E) \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \mathcal{O}_E \rightarrow 0.$$

By vanishing theorem  $H^1(\tilde{X}; \tilde{L}^\nu \otimes \mathcal{O}(-E)) = 0$  for  $\nu \gg 0$ .

Similarly:

c)  $\phi_{L^\nu}$  has nondegenerate differential at  $x$ . Equivalently

$$H^0(X; L^\nu \otimes m_x) \rightarrow L_x^\nu \otimes T_x^* X = H^0(X; L^\nu \otimes \Omega_X^1 \otimes \mathcal{O}_X/m_x)$$

is surjective.

c') the restriction

$$H^0(\tilde{X}; \tilde{L}^\nu \otimes I_E) \rightarrow H^0(E; \tilde{L}_{|E}^\nu \otimes I_E) = H^0(\tilde{X}; \tilde{L}^\nu \otimes \mathcal{O}_{\tilde{X}}/I_E \otimes I_E)$$

is surjective, where  $\tilde{X} = Bl_x X$ ,  $\tilde{L}$  where  $\tilde{X}$  is the pull-back of  $L$ ,  $E$  is the exceptional divisor,  $I_E$  is the ideal sheaf of  $E$ . Note that  $I_E$  restricted to  $E = \mathbb{P}(T_x X)$  is isomorphic to  $\mathcal{O}(1)$ , hence

$$H^0(E; \tilde{L}_{|E}^\nu \otimes I_E) = L_x^\nu \otimes T_x^* X.$$

The restriction map is a part of an exact sequence

$$\rightarrow H^0(\tilde{X}; \tilde{L}^\nu \otimes I_E) \rightarrow H^0(\tilde{X}; \tilde{L}^\nu \otimes \mathcal{O}_{\tilde{X}}/I_E \otimes I_E) \rightarrow H^1(\tilde{X}; \tilde{L}^\nu) \rightarrow$$

**13.12 Kähler manifolds with integral (up to a scalar) Kähler form are projective.** If  $(X, \omega)$  is a Kähler manifold and  $[\omega]$  is of the form  $\lambda[\omega']$  for  $\lambda \in \mathbb{R}_+$ ,  $[\omega'] \in H^*(X; \mathbb{Z})$ , then  $X$  embeds into a projective space.

- Proof. It is enough to show that: if  $[\omega]$  is integral, then there exist a holomorphic line bundle  $L$  with a curvature form equal to  $\frac{2\pi}{i}c_1(L)$ .
- The short exact sequence

$$\mathbb{Z} \rightarrow \mathbb{C} \rightarrow \mathcal{O}_X$$

of sheaves induces a long exact sequence

$$\rightarrow \text{Pic}(X) = H^1(X, \mathcal{O}_X^*) \xrightarrow{c_1} H^2(X; \mathbb{Z}) \xrightarrow{\iota} H^2(X; \mathcal{O}_X) \rightarrow .$$

Assume that  $[\omega']$  is integral. We will show that  $[\omega']$  lies in the image of  $c_1$ , or equivalently it belongs to the kernel of the map  $\iota$ . The map  $\iota$  factors as follows

$$H^2(X; \mathbb{Z}) \rightarrow H^2(X; \mathbb{C}) = H^{2,0}(X) \oplus H^{1,1}(X) \oplus H^{0,2}(X) \twoheadrightarrow H^{0,2}(X) = H^2(X; \mathcal{O}_X).$$

The second map is induced by the map of sheaves  $\mathbb{C} \rightarrow \mathcal{O}_X$ . The classes of the type  $(1, 1)$  lie in the kernel.

- It remains to show, that if  $c_1(L) = [\omega]$  then  $L$  admits a connection  $\nabla$  with

$$\frac{i}{2\pi}F_\nabla = \omega.$$

**13.13 adjusting the connection.** Suppose  $[\omega] = c_1(L) \in H^2(X)$ , where  $\omega$  is a real  $(1,1)$ -form. Then there exists a connection  $\nabla$  such that  $\omega = \frac{i}{2\pi}F_\nabla$ .

- Proof: locally in a trivialization  $F_\nabla = \bar{\partial}\partial \log(h(z))$ . For a different choice of a metric  $h' = e^\rho h$ . Hence

$$F_{\nabla'} = \bar{\partial}\partial \log(e^\rho h(z)) = F_\nabla + \bar{\partial}\partial \rho.$$

We want to find

$$F_{\nabla'} = -2\pi i \omega = F_\nabla + d\beta$$

for a given  $\beta \in A^1(X)$ . It remains to solve an equation

$$\bar{\partial}\partial \rho = d\beta.$$

Then  $h' = e^\rho h$  and  $\nabla'$  is the required connection.

- We apply  $\partial\bar{\partial}$ -lemma [Huy, Cor 3.2.10]: for a given exact form  $d\beta$  of the type  $(1,1)$ , there exists  $\rho$  such that  $d\beta = \bar{\partial}\partial \rho$ .
- Existence of  $\rho$  is the conclusion of  $\partial\bar{\partial}$ -lemma For  $\phi \in A^{p,q}$

$$\phi = d\beta \quad \Rightarrow \quad \exists \gamma \quad \phi = \partial\bar{\partial}\gamma$$

### 13.14 Numerical criteria for admitting a positive connection, i.e. ampleness

- Nakai-Moishezon criterion
- Kleiman criterion

[Ten temat już należy do innego przedmiotu.]

## 14 Dowód twierdzenia Kodairy-Nakano o znikaniu

**14.1** Jeśli  $L$  i  $K$  dodatnie, to  $L \otimes K$  też. Jeśli  $K = L^{\otimes n}$  jest dodatnie, to  $L$  też. Obcięcie zachowuje dodatniość.

Poniżej używamy  $E$  jako oznaczenie wiązki, bo  $L$  jest zarezerwowane na operator Lefschetza. Zakładamy, że  $X$  jest wartą rozmaitością analityczną.

**14.2** Ustalamy metrykę hermitowską na  $X$  i  $E$ . Definiujemy skręcone harmoniczne

$$\mathcal{H}^{p,q}(E) := \ker(\Delta_E) \subset A^{p,q}(E), \quad \Delta_E = \bar{\partial}_E \bar{\partial}_E^* + \bar{\partial}_E^* \bar{\partial}_E$$

**14.3** Jeśli  $X$  jest zwarta to

$$A^{p,q}(E) = \mathcal{H}^{p,q}(E) \oplus \text{im}(\bar{\partial}_E) + \text{im}(\bar{\partial}_E^*)$$

**14.4** Z rozkładu Hodge'a dla  $\bar{\partial}$

$$H^{p,q}(X; E) = \mathcal{H}^{p,q}(E) = \ker(\Delta_E).$$

**14.5** Mamy

- Operator  $L : A^{p,q}(X; E) \rightarrow A^{p+1,q+1}(X; E)$  i sprzężony  $L^*$
- $\nabla = \nabla^{1,0} + \bar{\partial}_E$  oraz tożsamość (u Huybrechtsa nazwana tożsamością Nakano)

$$[L^*, \bar{\partial}_E] = -i(\nabla^{1,0})^*,$$

która jest uogólnieniem tożsamości Kählera  $[L^*, \bar{\partial}] = -i\partial^*$ .

**14.6** (Znikanie Kodairy-Nakano) Jeśli  $E$  jest dodatnia, to  $H^{p,q}(X; E) = H^q(X; \Omega_X^p \otimes E) = 0$  dla  $p + q > \dim X$ .

Forma krzywizny  $F_{\nabla}$  jest typu  $(1,1)$ , lokalnie

$$F_{\nabla} = dA = \bar{\partial}\partial(\log h).$$

Lokalnie dla przekroju  $\eta$

$$\nabla^{1,0}\eta = \partial\eta + \partial(\log h) \wedge \eta$$

$$\bar{\partial}_E(\nabla^{1,0}\eta) = (-\partial\bar{\partial}\eta + \bar{\partial}(\partial(\log h) \wedge \eta)) = -\partial\bar{\partial}\eta - \partial(\log h) \wedge \bar{\partial}\eta + \bar{\partial}\partial(\log h) \wedge \eta$$

$$\nabla^{1,0}(\bar{\partial}_E\eta) = \partial\bar{\partial}\eta + \partial(\log h) \wedge \bar{\partial}\eta$$

$$\boxed{F_{\nabla} \wedge \eta = \bar{\partial}_E \nabla^{1,0}\eta + \nabla^{1,0} \bar{\partial}_E \eta}$$

Niech  $\eta \in \mathcal{H}^{p,q}(E)$ ,

$$\bar{\partial}_E \eta = 0, \quad \bar{\partial}_E^* \eta = 0$$

Wtedy

$$\boxed{F_{\nabla} \wedge \eta = \bar{\partial}_E \nabla^{1,0}\eta}$$

Stąd

$$\begin{aligned} i\langle L^* F_{\nabla} \eta, \eta \rangle &= i\langle L^* \bar{\partial}_E \nabla^{1,0}\eta, \eta \rangle \stackrel{Nakano}{=} i\langle (\bar{\partial}_E L^* - i(\nabla^{1,0})^*) \nabla^{1,0}\eta, \eta \rangle = \\ &= i\langle (\bar{\partial}_E L^*, \eta) + \langle (\nabla^{1,0})^* \nabla^{1,0}\eta, \eta \rangle = i\langle L^*, \partial_E^* \eta \rangle + \langle \nabla^{1,0}\eta, \nabla^{1,0}\eta \rangle = \langle \nabla^{1,0}\eta, \nabla^{1,0}\eta \rangle \geq 0 \end{aligned}$$

Podobnie

$$\begin{aligned} i\langle F_{\nabla} L^* \eta, \eta \rangle &= i\langle (\bar{\partial}_E \nabla^{1,0} + \nabla^{1,0} \bar{\partial}_E) L^* \eta, \eta \rangle = i\langle (\nabla^{1,0} \bar{\partial}_E) L^* \eta, \eta \rangle = i\langle \nabla^{1,0} (L^* \bar{\partial}_E + i(\nabla^{1,0})^*) \eta, \eta \rangle = \\ &= -\langle \nabla^{1,0} (\nabla^{1,0})^* \eta, \eta \rangle = -\langle (\nabla^{1,0})^* \eta, (\nabla^{1,0})^* \eta \rangle \leq 0 \end{aligned}$$

Stąd

$$i\langle [L^*, F_{\nabla}] \eta, \eta \rangle = \|(\nabla^{1,0})^* \eta\|^2 + \|\nabla^{1,0} \eta\|^2 \geq 0$$

Ale

$$iF_{\nabla} \wedge - = 2\pi L$$

bo  $L$  jest dodatnia, więc  $\frac{i}{2\pi} F_{\nabla}$  jest formą Kählera. Stąd

$$i\langle [L^*, F_{\nabla}] \eta, \eta \rangle = 2\pi \langle [L^*, L] \eta, \eta \rangle = -2\pi \langle H \eta, \eta \rangle = 2\pi(n - (p + q))\|\eta\|^2.$$

Jeśli  $n - (p + q) \leq 0$ , to  $i\langle [L^*, F_{\nabla}] \eta, \eta \rangle \leq 0$ . Zatem  $\|\eta\|^2 = 0$ .

**14.7** Wniosek: jeśli  $E^{\otimes n} = \mathcal{O}(1)_X$  dla pewnego zanurzenia  $X \subset \mathbb{P}^N$ , to  $H^k(X, \Omega_X^n \otimes E) = 0$  dla  $k > 0$ .

**14.8** Poprzez dualność Serre'a (lub bezpośrednio, korzystając z tego, że  $\frac{i}{2\pi} F_{\nabla_{E^*}} = -\omega$ ):

$$H^k(X, E^*) = 0$$

dla  $k < n$ .

# Complex Manifolds - problem list no. 1 - zadania na 17.10


1 Show that any compact complex curve of genus 1 is isomorphic with  $\mathbb{C}/\langle 1, \tau \rangle$ .

Hint: Show that  $\mathbb{Z}^2$  is not isomorphic with a discrete subgroup of  $SL_2(\mathbb{R})$ .

2 Show that the following map is a bijection:

$$\begin{aligned} \mathbb{H}/PSL_2(\mathbb{Z}) &\longrightarrow \{\text{isomorphism classes of curves of genus 1}\} \\ \tau &\longmapsto \mathbb{C}/\langle 1, \tau \rangle. \end{aligned}$$

3 What are the stabilizers of points in  $\mathbb{H}$  with respect to the action of  $PSL_2(\mathbb{Z})$ ? Describe the quotient space  $\mathbb{H}/PSL_2(\mathbb{Z})$ . Is it homeomorphic to some surface?



4 Quadrics in  $\mathbb{P}^n$  for  $n \leq 5$ :

a) Show that Plücker embedding  $Gr(2, 4) \hookrightarrow \mathbb{P}(\Lambda^2 \mathbb{C}^4)$  is a quadric.

b) Show that the image of the Veronese embedding  $\mathbb{P}^1 \hookrightarrow \mathbb{P}^2$  is a quadric.

c) Show that the image of the Segre embedding  $\mathbb{P}^1 \times \mathbb{P}^1 \hookrightarrow \mathbb{P}^3$  is a quadric.

d) Can you say anything particular about the quadric in  $\mathbb{P}^4$ ? (keyword: Lagrangian Grassmannian)

*Check the underlined words e.g. in Wikipedia.*

5 [Referat dla KP] Let  $\Lambda$  be a lattice in  $\mathbb{C}$ . Let

$$\wp(z) = z^{-2} + \sum_{w \in \Lambda \setminus \{0\}} ((z - w)^{-2} - w^{-2})$$

be the Weierstrass function. (Argue that the sum is convergent and differentiable term by term.) Show that  $\wp(z)$  is  $\Lambda$ -periodic. Moreover

$$\wp'(z)^2 = 4\wp(z)^3 - g_2\wp(z) - g_3,$$

where

$$g_2 = 60 \sum_{w \in \Lambda \setminus \{0\}} w^{-4},$$

$$g_3 = 140 \sum_{w \in \Lambda \setminus \{0\}} w^{-6}.$$

Show that

$$z \mapsto [\wp(z) : \wp'(z) : 1]$$

defines a continuous map  $\mathbb{C}/\Lambda \rightarrow \mathbb{P}^2$ , an isomorphism on the image given by the equation

$$Q(x, y, z) = y^2z - 4x^3 + g_2x + g_3 = 0.$$

(see Kirwan, *Complex Algebraic Curves*, Chapter 5).



## Complex Manifolds - problem list no. 2 - zadania na 31.10

**1** Suppose  $n > 1$ ,  $f : \mathbb{C}^n \rightarrow \mathbb{C}$  holomorphic. Show that if the zero set  $Z(f)$  is compact, then it is empty.

**2** Suppose  $U \subset \mathbb{C}^n$  is open and non-empty. Is the ring of holomorphic functions on  $U$  a unique factorisation domain?

**3** Find the Weierstrass polynomial for  $f(z_1, z_2) = z_1^3 z_2 + z_1 z_2 + z_1^2 z_2^2 + z_2^2 + z_1 z_2^3$  at 0.

**4** Suppose  $U \subset \mathbb{C}^n$  is connected and  $0 \neq f : U \rightarrow \mathbb{C}$  is holomorphic. Show that  $U \setminus Z(f)$  open dense and connected.

**5** [Referat dla MJ] Let  $f : U \rightarrow V$  be a holomorphic function, which is bijective. Show, that  $f^{-1}$  is holomorphic. [Huybrechts: Proposition 1.1.13]

## Complex Manifolds - problem list no. 3 - zadania na 7.10

**1** *Zadanie 4 z listy nr 2.*

**2** Deduce formally the implicit function theorem (for  $f : \mathbb{C}^n \rightarrow \mathbb{C}$  with  $\frac{\partial f}{\partial z_1} \neq 0$ ) as a special case of Weierstrass theorem.

**3** *Lagrangian Grassmannian  $LG_2$ .* Consider the standard symplectic form  $\omega$  in  $\mathbb{C}^4$ . It is given in the standard coordinates by the matrix

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}.$$

Show that the set of planes  $L \subset \mathbb{C}^4$  satisfying the condition  $\omega|_L \equiv 0$ , considered as a subset of the Grassmannian  $Gr_2(\mathbb{C}^4)$  is a complex submanifold. It can be embedded into  $\mathbb{P}^4$  as a nonsingular quadric.

(Symplectic = antisymmetric, nondegenerate 2-form.)

**4** *Cohomology groups of complex manifolds with coefficients in  $\mathbb{Z}$  or de Rham cohomology*

- a) Describe the cohomology ring of the complex projective space  $\mathbb{P}^n$ .
- b) Compute cohomology groups of the Grassmannian  $Gr_2(\mathbb{C}^4)$ .
- c) Compute  $H^*(LG_2)$ .

Hint: show that there exist decompositions into cells isomorphic to complex affine spaces.

**5** Jeśli starczy czasu, możemy przedyskutować zadanie 1.1.11 z Huybrechtsa (str. 19).

## Complex Manifolds - problem list no. 4 - zadania na 14.11

**1** Niech  $Q$  będzie zespoloną formą dwuliniową na  $\mathbb{C}^n$ . Utożsamiamy  $\mathbb{C}^n$  z  $\mathbb{R}^{2n}$ . Udowodnić, że jeśli  $v \in \mathbb{R}^{2n}$  jest wektorem własnym części rzeczywistej  $\operatorname{Re} Q$  o wartości własnej  $\lambda$  to  $iv$  jest wektorem własnym dla wartości  $-\lambda$ .

(Uwaga: wektor  $v$  jest wektorem własnym formy dwuliniowej  $B$  jeśli dla każdego  $w$  mamy  $B(v, w) = \lambda \langle v, w \rangle$ . W naszym przypadku zakładamy, że iloczyn skalarny jest niezmienniczy ze względu na mnożenie przez  $i$ , tzn  $\langle iv, iw \rangle = \langle v, w \rangle$  )

**2** Niech  $X = \{z \in \mathbb{C}^n : \sum_{k=1}^n z_k^2 = 1\}$ . Wykazać, że zbiór  $X$  jest homotopijnie równoważny ze sferą  $S^{n-1}$ . (A nawet jest homeomorficzny z przestrzenią styczną  $TS^{n-1}$ .)

**3** Niech  $X = \{(x, y) \in \mathbb{C}^2 : 2x^3 - 6x - 3y^2 = 0\}$  oraz  $f : X \rightarrow \mathbb{R}$ ,  $f(x, y) = \operatorname{Re} y$ . Czy punkty krytyczne  $f$  są niezdegenerowane (tzn czy jeśli  $D(f) = 0$  to  $D^2(f)$  jest niezdegenerowana)? Jakie są indeksy?

**4** Niech  $f : \mathbb{P}^n \rightarrow \mathbb{R}$  będzie zadane formułą

$$f([z_0 : z_1 : \cdots : z_n]) = \sum_{k=0}^n k \frac{|z_k|^2}{||z||^2}.$$

Wykazać, że  $f$  ma niezdegenerowane punkty krytyczne i obliczyć ich indeksy.

**5** Referat dla S.S.: Opisać rozkład grassmanianu na komórki algebraiczne

$$Gr_k(\mathbb{C}^n) = \bigsqcup_{\lambda \in ??} \mathbb{C}^{n_\lambda}.$$

[Griffiths-Harris, Principles of Algebraic Geometry, roz I.5 str 193-7]

## Complex Manifolds - problem list no. 5 - zadania na 21.11

**1** Niech  $X = \{(x, y) \in \mathbb{C}^2 : 2x^3 - 6x - 3y^2 = 0\}$  oraz  $f : X \rightarrow \mathbb{R}$ ,  $f(x, y) = \operatorname{Re} y$ . Czy punkty krytyczne  $f$  są niezdegenerowane (tzn czy jeśli  $D(f) = 0$  to  $D^2(f)$  jest niezdegenerowana)? Jakiekolwiek są indeksy?

**2** Show that Hodge  $*$  satisfies

- (i)  $*^2 = (-1)^{k(d-k)}$  for  $k$ -forms,
- (ii)  $\langle \alpha, *\beta \rangle = (-1)^{k(d-k)} \langle *\alpha, \beta \rangle$ .

**3** Let  $M = S^1$ .

- Find the formula for  $d^*$  for 1-forms.
- Find the formula for  $\Delta$  for 0-forms and 1-forms.
- Find all eigenvalues of  $\Delta$  acting on  $\mathcal{A}^k(S^1)$  for  $k = 0, 1$ .
- Solve the heat equation for  $\alpha : \mathbb{R} \rightarrow \mathcal{A}^k(S^1)$ :

$$\frac{d}{dt}\alpha_t = -\Delta\alpha_t$$

with the initial condition

$$\alpha_0 = \left( a + \sum_{k=1}^{\infty} b_k \sin(kt) + c_k \cos(kt) \right) dt.$$

**4** Compute Laplacian for the torus  $\mathbb{R}^n/\mathbb{Z}^n$  with the standard (flat) metric. Find the spaces of harmonic forms.

**5** (Referat dla ML) Let  $\alpha_t$  be a solution of the heat equation

$$\frac{d}{dt}\alpha_t = -\Delta\alpha_t$$

with the initial condition  $\alpha_0$ . Show that if  $d\alpha_0 = 0$ , then  $d\alpha_t = 0$  and the cohomology class  $[\alpha_t] \in H^*(M)$  for  $t \geq 0$  is constant. The limit is equal to the harmonic representative of  $[\alpha_t]$ .

[Donu Arapura, Algebraic Geometry over the Complex Numbers, Universitext 2012, Section 8.3]

## Complex Manifolds - problem list no. 6 - zadania na 28.11

**1** Show that Hodge  $*$  satisfies

- (i)  $*^2 = (-1)^{k(d-k)}$  for  $k$ -forms,
- (ii)  $\langle \alpha, *\beta \rangle = (-1)^{k(d-k)} \langle *\alpha, \beta \rangle$ .

**2** Let  $M = S^1 = \mathbb{R}/2\pi\mathbb{Z}$ . Solve the heat equation for  $\alpha : \mathbb{R} \rightarrow \mathcal{A}^k(S^1)$ :

$$\frac{d}{dt}\alpha_t = -\Delta\alpha_t$$

with the initial condition

$$\alpha_0 = \delta_0 dt,$$

where  $\delta_0$  is the Dirac delta. Show that for  $t > 0$  the form  $\alpha_t$  is smooth.

Uwaga: rozważamy tu formy o współczynnikach, które są dystrybucjami. (Takie formy nazywamy prądami, ang. „currents”.) Dla  $n$ -wymiarowej rozmaitości takie  $k$ -formy działają na  $(n-k)$ -formach próbnych o zwartym nośniku w sposób oczywisty (?). Rozwinięcie dystrybucji w szereg Fouriera traktujemy czysto formalnie.

**3** Compute Laplasian for the torus  $(S^1)^n$  with the standard (flat) metric. Find the spaces of harmonic forms.

**4** Let  $I$  be a standard complex structure in  $\mathbb{C}^n = \mathbb{R}^n \oplus i\mathbb{R}^n \simeq \mathbb{R}^{2n}$ . Consider the following subgroups of  $GL_{2n}(\mathbb{R})$ :

- $GL_n(\mathbb{C})$  consisting of  $A$  such that  $AI = IA$ ,
- $Sp_n(\mathbb{R})$  consisting of  $A$  such that  $A^T I A = I$ ,
- $O(2n)$  consisting of  $A$  such that  $A^T A = id$ .

Show that intersections of any two of three groups above is equal to  $U_n \subset GL_n(\mathbb{C})$ , the group of unitary matrices, i.e. the subgroup of  $GL_n(\mathbb{C})$  preserving the hermitian product.

**5** Let  $V$  be a complex vector space with a hermitian structure. Consider the space  $\Lambda^\bullet = \Lambda^\bullet(V^* \otimes_{\mathbb{R}} \mathbb{C})$  with operations  $L, H, L^*$ :

- $L(\alpha) = \omega \wedge \alpha$ ,
- $H|_{\Lambda^k} = \text{multiplication by } k - n$ ,
- $L^*(\alpha) = (-1)^k * L(*\alpha)$  for  $\alpha \in \Lambda^k$  (show that indeed this is the adjoint operator to  $L$ ).

Show that

- $[L, L^*] = H$ ,
- $[H, L] = 2L$ ,
- $[H, L^*] = -2L^*$ .

That is, show that  $\Lambda^*$  is a representation of  $\mathfrak{sl}_2(\mathbb{Z})$ .

Hint: Check it for  $\dim_{\mathbb{C}} V = 1$  and proceed by induction.

**6** Decompose the representation  $\Lambda^\bullet(V^* \otimes_{\mathbb{R}} \mathbb{C})$  into irreducible representations assuming that  $\dim V = 1, 2$  or (maybe)  $3$ .

## Complex Manifolds - problem list no. 7 - zadania na 5.12

**1** Decompose the representation  $\Lambda^\bullet(V^* \otimes_{\mathbb{R}} \mathbb{C})$  into irreducible representations assuming that  $\dim V = 1, 2$  or (maybe)  $3$ .

**2** [Fubini-Study metric on  $\mathbb{P}^n$ ] Consider a local holomorphic section of the quotient bundle  $\mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{P}^n$

$$s : \mathbb{P}^n \supset U \rightarrow \mathbb{C}^{n+1} \setminus \{0\}$$

Show that the form

$$\omega = \frac{i}{2\pi} \partial \bar{\partial} \log(|s(z)|^2)$$

Does not depend on the choice of  $\alpha(z)$ . Hence can be glued to a global form on  $\mathbb{P}^n$ . In particular, in local coordinates on  $U_0 = \{z_0 \neq 0\}$

$$\omega = \frac{i}{2\pi} \partial \bar{\partial} \log(1 + \sum_{k=1}^n |z_k|^2).$$

Show that the resulting form is the Fubini-Study form.

Hint: Perform the calculation for  $s(z) = \alpha(z)(1, z_1, \dots, z_n)$ .

**3** Prove the second Hodge identity  $[L^*, \partial] = i\bar{\partial}^*$  for the flat metric: We decompose  $\bar{\partial} = \sum \bar{\partial}_k$  and  $L^* = \sum L_k^*$ . Show that  $\bar{\partial}_k^* = -2\iota_{\bar{v}_k} \frac{\partial}{\partial z_k}$ , where  $\bar{v}_k = \frac{\partial}{\partial \bar{z}_k}$ . Note  $\partial_\ell$  commutes with  $L_k^*$  for  $k \neq \ell$ . It remains to check  $[L_k^*, \partial_k]$  for  $\alpha = f dz_I \wedge d\bar{z}_J$ , considering 4 cases  $k \in$  or  $\notin$  to  $I$  and  $J$ . For example: suppose  $k \in I, k \in J$ . That is  $I = \{k\} \cup I', J = \{k\} \cup J'$ :

$$\begin{aligned} [L_k^*, \partial_k] f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'} &= \\ L_k^* \partial_k (f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'}) - \partial_k L_k^* (f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'}) &= \\ 2i \partial_k (f dz_{I'} \wedge d\bar{z}_{J'}) &= \\ 2i \frac{\partial f}{\partial z_k} dz_k \wedge dz_{I'} \wedge d\bar{z}_{J'} &= \\ 2i \frac{\partial f}{\partial z_k} \iota_{\bar{v}_k} (d\bar{z}_k \wedge dz_k \wedge dz_{I'} \wedge d\bar{z}_{J'}) &= \\ i \bar{\partial}^* (f dz_k \wedge d\bar{z}_k \wedge dz_{I'} \wedge d\bar{z}_{J'}), \end{aligned}$$

Compute another case, e.g.  $k \notin I, k \in J$ .

**4** Referat dla BS: Holomorphic Poincaré Lemma (Huybrechts 1.3.7&8).

## Complex Manifolds - problem list no. 8 - zadania na 12.12

**1** Compute the Dolbeault cohomology  $H^q(M; \Omega_M^p) = H^q(A^{p,\bullet}(M), \bar{\partial})$  for  $M = \mathbb{C} \setminus \{0\}$ .

**2** Let  $\mathbb{D}$  be the unit disc. Suppose  $\mathcal{X} \subset \mathbb{P}^n \times \mathbb{D}$  is a submanifold, and  $X_t = \mathcal{X} \cap (\mathbb{P}^n \times \{t\})$  is smooth for each  $t \in \mathbb{D}$ . We assume that for each  $t$  the metric on  $X_t$  is induced from the standard Fubini-Study metric on  $\mathbb{P}^n$ . Show that the volume  $\text{vol}(X_t)$  is constant.

*A shorter formulation: the volume is constant in families of submanifolds.*

**3** Let  $N \subset M$  be a submanifold of a compact Kähler manifold of codimension  $c$ . Show that the (Poincaré dual of) the fundamental class  $[N] \in H^{2c}(M; \mathbb{C})$  is nonzero and of the type  $(c, c)$ .

**4** Define the set  $\mathcal{K}_X \subset H^{1,1}(X) \cap H^2(X; \mathbb{R})$  consisting of the classes  $[\omega]$  such that  $\omega$  is the minus imaginary part of a Kähler metric. Show that  $\mathcal{K}_X$  is an open conical set, not containing any affine line.

**5** Let  $M = \text{Bl}_x \mathbb{P}^2$  be the blow-up of the projective plane, see [Huybrechts §2.5]. Compute Euler characteristic and the intersection form on  $H^2(M)$ .

(Alternativ description:  $M$  is homeomorphic to the connected sum

$$\mathbb{P}^2 \# \overline{\mathbb{P}^2} := (\mathbb{P}^2 \setminus B_\epsilon) \sqcup (\mathbb{P}^2 \setminus B_\epsilon) / \sim,$$

where we glue the points of the boundaries of the balls  $B_\epsilon$ , so that the orientations of  $\mathbb{P}^2$ 's do not agree.)

**6** Refrat KP: *Kähler potential*. Let  $M$  be a hermitian manifold (i.e. a complex manifold with  $I$ -invariant Riemannian structure). Show that the 2-form  $\omega(v, w) = \langle I(v), w \rangle$  is closed if and only if locally  $\omega = i\partial\bar{\partial}f$  for a smooth function  $f : M \rightarrow \mathbb{R}$ .

## Complex Manifolds - problem list no. 9 - zadania na 19.12

**1** Compute the Dolbeault cohomology  $H^q(M; \Omega_M^p) = H^q(A^{p,\bullet}(M), \bar{\partial})$  for  $M = \mathbb{C} \setminus \{0\}$ .

*Wskazówka: Zostało do obliczenia  $H^1(M, \mathcal{O}_M) \simeq H^1(M, \Omega_M^1)$ . Powinno wyjść 0, czyli trzeba wykazać, że każda forma  $f(z)d\bar{z}$  jest różniczką. Proponuję najpierw sprawdzić czy jest to prawda dla  $M = \mathbb{C}$  a potem wykorzystać  $\exp : \mathbb{C} \rightarrow \mathbb{C} \setminus \{0\}$ .*

**2** Define the set  $\mathcal{K}_X \subset H^{1,1}(X) \cap H^2(X; \mathbb{R})$  consisting of the classes  $[\omega]$  such that  $\omega$  is the minus imaginary part of a Kähler metric. Show that  $\mathcal{K}_X$  is an open conical set, not containing any affine line.

**3** Let  $M$  be the connected sum

$$\mathbb{P}^2 \# \mathbb{P}^2 := (\mathbb{P}^2 \setminus B_\epsilon) \sqcup (\mathbb{P}^2 \setminus B_\epsilon) / \sim,$$

where we glue the points of the boundaries of the balls  $B_\epsilon$ , so that the orientations of  $\mathbb{P}^2$ 's agree.

a) Compute the Betti numbers (i.e. the dimensions  $\dim H^k(M)$ ) and the signature of  $M$ .

b) Using the formula relating Hodge numbers with the signature deduce that  $M$  does not admit any structure of Kähler manifold.

**4** Compute Čech cohomology  $\check{H}^1(\mathcal{U}; \Omega^1)$ , where  $\mathcal{U} = \{U_0, U_1\}$  is the standard cover of  $\mathbb{P}^1$ ,  $U_i \simeq \mathbb{C}$  for  $i = 0, 1$ .

**5** (*Trudniejsze*) Construct a lattice  $\Lambda \subset \mathbb{C}^2$  such that  $\mathbb{C}^2/\Lambda$  cannot be embedded into a projective space.

*Wskazówka: Jeśli  $M$  można zanurzyć w  $\mathbb{P}^n$ , to  $\omega_{FS}$  pochodzi z  $H^2(M; \mathbb{Z})$ . Skonstruować taką kratę, by  $H^2(M; \mathbb{Q}) \cap H^{1,1}(M) = 0$  (jako podprzestrzeń  $H^2(M; \mathbb{C})$ ).*

**6** Referat S.S.: Let  $M = Bl_x \mathbb{P}^2$  be the blow-up of the projective plane, see [Huybrechts §2.5]. Compute Euler characteristic and the intersection form on  $H^2(M)$ .

(Alternativ description:  $M$  is homeomorphic to the connected sum

$$\mathbb{P}^2 \# \overline{\mathbb{P}^2} := (\mathbb{P}^2 \setminus B_\epsilon) \sqcup (\mathbb{P}^2 \setminus B_\epsilon) / \sim,$$

where we glue the points of the boundaries of the balls  $B_\epsilon$ , so that the orientations of  $\mathbb{P}^2$ 's do not agree.)



## Complex Manifolds - problem list no. 10 - zadania na 9.01.2023

**1** Construct a lattice  $\Lambda \subset \mathbb{C}^2$  such that  $\mathbb{C}^2/\Lambda$  cannot be embedded into a projective space.

*Wskazówka: Jeśli  $M$  można zanurzyć w  $\mathbb{P}^n$ , to  $\omega_{FS|_M}$  pochodzi z  $H^2(M; \mathbb{Z})$ . Skonstruować taką kratę, by  $H^2(M; \mathbb{Z}) \cap H^{1,1}(M) = 0$ . Dla wygody  $\mathbb{Z}$  można zastąpić przez  $\mathbb{Q}$ . Rozważamy  $H^2(M; \mathbb{Z})$ ,  $H^2(M; \mathbb{Q})$ ,  $H^2(M; \mathbb{R})$  jako podgrupy  $H^2(M; \mathbb{C})$ .*

*Dokładniejsze wskazówki: Zapiszmy  $\beta = \frac{i}{2} \sum_{i,j} a_{i,j} dz_i \wedge d\bar{z}_j$ ,  $a_{i,j} \in \mathbb{C}$ . To jest ogólna postać klasy kokohomologii  $[\beta] \in H^{1,1}(M) \subset H^2(M; \mathbb{C})$  (ta podprzestrzeń nie zależy od wyboru formy Kählera).*

a) Dla jakich  $a_{i,j}$  klasa  $[\beta] \in H^2(M; \mathbb{R})$ ?

b) Można założyć, że krata jest rozpięta przez wektory  $\alpha_1 = (1, 0)$ ,  $\alpha_2 = (0, 1)$  oraz  $\alpha_3$  i  $\alpha_4 \in \mathbb{C}^2$ . Niech  $T_{k,\ell} = (\mathbb{R}\alpha_k + \mathbb{R}\alpha_\ell)/(\mathbb{Z}\alpha_k + \mathbb{Z}\alpha_\ell)$ . Obliczyć całki  $\int_{T_{k,\ell}} \beta$ .

c) Zobaczyć, że dla ogólnego wyboru wektorów  $\alpha_3$  i  $\alpha_4$  nie można dobrać współczynników  $a_{i,j}$  tak by wszystkie całki  $\int_{T_{k,\ell}} \beta$  były wymierne.

**2** Compute  $H^1(\mathbb{P}^n; \mathcal{O}_{\mathbb{P}^n}^*)$  which parametrizes line bundles over  $\mathbb{P}^n$ . (Hint: Use the exponential exact sequence.)

**3** Let  $X = Bl_{pt} \mathbb{P}^2$ .

a) Construct a Kähler form on  $X$ .

b) Find an embedding of  $X$  into a projective space.

c) Find all the cohomology classes  $[\alpha] \in H^{1,1}(X)$  satisfying  $\int_C \alpha > 0$  for all complex curves  $C \subset X$  and  $\int_X \alpha^2 > 0$ . (It is enough to consider  $C = E$  – exceptional divisor or  $C = H$  – the pullback of a line in  $\mathbb{P}^2$ .)

Remark: See also *Nakai–Moishezon criterion*.

Podobne pytania można zadać dla  $X = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(k))$ , tj dla powierzchni Hirzebrucha. Jeśli zostanie czas, to zajmiemy się tym.

**4** Let  $E$  and  $F$  be complex vector bundles over a manifold. Let  $\nabla_E$  and  $\nabla_F$  be their connections. Construct a connection on the vector bundle  $\underline{\text{Hom}}(E, F)$ .

**5** Referat MJ: Opowiedzieć o dualności Serra [Huybrechts: Prop. 4.1.15]

## Complex Manifolds - problem list no. 11 - zadania na 16.01.2023

1 Definiujemy wielomiany  $\sigma_n : \text{Macierze}(n \times n) \rightarrow \mathbb{C}$  wzorem

$$\sum \sigma_k(X) t^{n-k} = \det(X + tI).$$

Wykazać, że każdy Ad-niezmieniczny wielomian na przestrzeni macierzy można wyrazić za pomocą wielomianów  $\sigma_n$ .

2 Niech  $E$  będzie wiązką zespoloną rangi  $n$  (nad dowolną bazą).

(a) Udowodnić  $c_1(E) = c_1(\Lambda^n E)$ .

(b) Załóżmy, że  $n = 3$ , wyrazić klasy Cherna wiązki  $\Lambda^2 E$  za pomocą klas Cherna  $E$ .

Wskazówka: Zastosować zasadę rozszczepienia i wzór Whitneya.

3 Niech  $E$  będzie wiązką holomorficzną nad bazą, która jest rozmaitością zespoloną,  $rk(E) = r$ . Niech  $s : X \rightarrow E$  będzie holomorficznym przekrojem wiązki, oraz  $Z(s) \subset X$  zbiorem jego zer.

(a) Zakładając, że  $s$  jest transwersalny do przekroju zerowego udowodnić, że klasa Poincaré dualna do  $[Z(s)]$  jest równa najwyższej klasie Cherna (tzn  $c_r(E)$ ). Oznacza to, że dla każdej formy  $\alpha$  o zwartym nośniku

$$\int_X \alpha \cdot c_r(E) = \int_{Z(s)} \alpha.$$

(b) Uogólnić powyższe stwierdzenie, przy założeniu, że  $s$  jest generycznie transwersalne do przekroju zerowego, tzn zbiór punktów nietranswersalności  $\Sigma(s)$  jest mniejszego wymiaru niż  $Z(s)$ .

Wskazówka: Zobaczyć, że w interesującej nas gradacji  $H^{2\text{codim}Z(s)}(X) \simeq H^{2\text{codim}Z(s)}(X \setminus \Sigma(s))$ .

4 Wykazać, że jeśli wiązka  $E$  ma nigdzie nie zerujący się przekrój, to  $c_r(E) = 0$ . (Bez żadnych założeń o bazie).

5 Referat: Klasa Atiyaha [Huybrechts od 4.2.17 do Prop 4.2.19].

## Complex Manifolds - problem list no. 12 - zadania na 23.01.2023

**1** Wykazać, że jeśli wiązka  $E$ ,  $\text{rk}(E) = r$  ma nigdzie nie zerujący się przekrój, to  $c_r(E) = 0$ . (O bazie  $X$  zakładamy tylko, że jest przestrzenią topologiczną).

**2** Let  $E \rightarrow M$  be a bundle with connection,  $U \subset M$  is an open subset. Suppose we have two sets of sections trivializing  $E|_U$ :  $s_1, s_2, \dots, s_n$  and  $s'_1, s'_2, \dots, s'_n$ . Let  $G \in GL_n(C^\infty(U))$  be the transition matrix. Suppose that

$$\nabla = d + A$$

with respect to the first trivialization. Find the formula for the connection form  $A'$  with respect to the second trivialization.

**3** Construct the Euler exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n} \rightarrow \mathcal{O}(1)^{\oplus n+1} \rightarrow T\mathbb{P}^n \rightarrow 0.$$

Hint: Let  $p : \mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{P}^n$  be the projection. For any  $v \in \mathbb{C}^{n+1} \setminus \{0\}$  we have a surjection

$$\mathbb{C}^{n+1} \simeq T_v \mathbb{C}^{n+1} \xrightarrow{T_v p} T_{[v]} \mathbb{P}^n.$$

Show that in fact we obtain a well defined surjection

$$\mathbb{C}^{n+1} \otimes \mathcal{O}(1) \rightarrow T\mathbb{P}^n.$$

**4** Let  $X \subset \mathbb{P}^n$  be a smooth hypersurface of degree  $d$ .

a) Compute  $\chi(X; \mathcal{O}(k))$  for  $n = 3$ ,  $d = 4$ ,  $k = 7$ .

b) Compute the Hodge numbers  $\dim H^q(X; \Omega^p)$  for  $n = 3$ ,  $d = 4$ .

Hint a): Let  $\mathcal{O}_X(d) := \mathcal{O}_{\mathbb{P}^n}(d)|_X$ . Then

$$TX \oplus \mathcal{O}_X(d)|_X \simeq T\mathbb{P}^n|_X$$

Hint b): let

$$\lambda_y(X) = \sum_{p=0}^{\dim X} \Lambda^p T^* X y^p$$

be a formal sum ( $y$  is a formal variable). Show that

$$\lambda_y(X)(\mathcal{O}_X + y\mathcal{O}_X(d)) = \lambda_y(\mathbb{P}^n).$$

**5** Referat ML: klasyczne twierdzenie Riemanna-Rocha dla krzywych i powierzchni, [Huybrechts §5.1, przykład 5.1.2i-ii].

## Tematy do opracowania:

- 1 Automorfizmy kuli w  $C^n$  i polidysku. Dlaczego kula nie jest biholomorficznie równoważna z polidyskiem? [Sha92]
- 2 Operator Laplasa na  $S^2 = \mathbb{P}^1$ , jego wartości i przestrzenie własne. Np [CH53, VII.7]
- 3 Kohomologiczne wnioski z rozkładu Białynickiego-Biruli. Omówić oryginalny dowód rozkładu kohomologii [Car02, §4.2] lub podać inny, np w oparciu o [Bro05].
- 4 Zbiory Steina (holomrficzne wypukłe) i ich kohomologie. Twierdzenia A i B Cartana o znikaniu wyższych kohomologii. Np [GR65]
- 5 Algebraiczne formy różniczkowe i kohomologie de Rhama - porównanie z topologicznymi dla algebraicznych rozmaitości afinicznych. [Gro66]
- 6 Rozwłóknienie Milnora dla izolowanych osobliwych punktów odwzorowania  $\mathbb{C}^n \rightarrow \mathbb{C}$ . [Mil68]
- 7 Teoria Picarda Lefschetza. [Ara12], [Ż06]
- 8 Teoria kobordyzmów i rozmaitości zespolone. Czy każda zorintowana rozmaitość jest kobordyczna z produktem  $\mathbb{P}^n$ 'ów? [MS74], [HBJ92]
- 9 Porównanie różnych definicji klas Cherna:  $c_1$  i wyższe klasy: definicja aksomatyczna, różniczkowa, przez teorię przeszkód, snopowo (dla  $c_1$ ). [MS74], [Huy05, §4.4], [FF16, §19.3].
- 10 Twierdzenia o znikaniu kohomologii dla dodatnich wiązek wektorowych (nie koniecznie dla wiązek liniowych). [Laz04, §7.3]
- 11 Opisać strukturę Hodge'a w kohomologiach powierzchni K3. [BHPVdV04, §VIII.3]. [Barth-Hulek-Chris-Van de Ven, Compact complex surfaces,

## References

- [Ara12] Donu Arapura. *Algebraic geometry over the complex numbers*. Universitext. Springer, New York, 2012.
- [BHPVdV04] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 2004.
- [Bro05] Patrick Brosnan. On motivic decompositions arising from the method of Białynicki-Birula. *Invent. Math.*, 161(1):91–111, 2005.
- [Car02] James B. Carrell. Torus actions and cohomology. In *Algebraic quotients. Torus actions and cohomology. The adjoint representation and the adjoint action*, volume 131 of *Encyclopaedia Math. Sci.*, pages 83–158. Springer, Berlin, 2002.
- [CH53] R. Courant and D. Hilbert. *Methods of mathematical physics. Vol. I*. Interscience Publishers, Inc., New York, N.Y., 1953.
- [FF16] Anatoly Fomenko and Dmitry Fuchs. *Homotopical topology*, volume 273 of *Graduate Texts in Mathematics*. Springer, [Cham], second edition, 2016.
- [GR65] Robert C. Gunning and Hugo Rossi. *Analytic functions of several complex variables*. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1965.

- [Gro66] A. Grothendieck. On the de Rham cohomology of algebraic varieties. *Inst. Hautes Études Sci. Publ. Math.*, (29):95–103, 1966.
- [HBJ92] Friedrich Hirzebruch, Thomas Berger, and Rainer Jung. *Manifolds and modular forms*. Aspects of Mathematics, E20. Friedr. Vieweg & Sohn, Braunschweig, 1992. With appendices by Nils-Peter Skoruppa and by Paul Baum.
- [Huy05] Daniel Huybrechts. *Complex geometry*. Universitext. Springer-Verlag, Berlin, 2005. An introduction.
- [Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. II*, volume 49 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2004. Positivity for vector bundles, and multiplier ideals.
- [Mil68] John Milnor. *Singular points of complex hypersurfaces*. Annals of Mathematics Studies, No. 61. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1968.
- [MS74] John W. Milnor and James D. Stasheff. *Characteristic classes*. Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974. Annals of Mathematics Studies, No. 76.
- [Sha92] B. V. Shabat. *Introduction to complex analysis. Part II*, volume 110 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1992. Functions of several variables, Translated from the third (1985) Russian edition by J. S. Joel.
- [Ż06] Henryk Żołądek. *The monodromy group*, volume 67 of *Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) [Mathematics Institute of the Polish Academy of Sciences. Mathematical Monographs (New Series)]*. Birkhäuser Verlag, Basel, 2006.

## Oral Exam – The List of Questions

- 1) Weierstrass preparation lemma
- 2) Hartogs theorem
- 3) Properties of the local ring of analytic functions  $\mathcal{O}_{\mathbb{C}^n,0}$
- 4) Classification of complex compact curves
- 5) Examples of complex compact manifolds. All about  $\mathbb{P}^n$ ,
- 6) Lefschetz hyperplane theorem (by Morse theory), cohomology of hypersurfaces in  $\mathbb{P}^n$
- 7) Hodge decomposition of  $C^\infty$  forms for compact Riemannian manifolds
- 8) Laplasian for compact Riemannian manifolds
- 9) Differential forms on complex manifolds, differentials  $\partial$  and  $\bar{\partial}$
- 10) Cousin problems and Čech cohomology
- 11) Dolbeault cohomology and their relation with  $H^*(X; \mathbb{C})$
- 12) Definition(s) of Kähler structure
- 13) Kähler identities
- 14) Hard Lefschetz theorem
- 15) The action of  $\mathfrak{sl}_2(\mathbb{Z})$  on differential forms and on cohomology of Kähler manifolds
- 16) Dualities: Poincaré, Serre, conjugation and Hodge diamond
- 17) Signature of Kähler manifolds
- 18) Fubini-Study metric on complex projective space
- 19) Connection and curvature for complex vector bundles,
- 20) Concordance of connection with complex and Hermitian structure
- 21) The first Chern class of a line bundle
- 22) Chern classes of holomorphic vector bundles
- 23) Hirzebruch-Riemann-Roch
- 24) Positive line bundles, Kodaira theorems